



Published in final edited form as:

Nat Rev Genet. 2015 January ; 16(1): 45–56. doi:10.1038/nrg3849.

Cystic fibrosis genetics: from molecular understanding to clinical application

Garry R. Cutting

McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University School of Medicine, 733 North Broadway, MRB 559, Baltimore, Maryland 21205, USA

Abstract

The availability of the human genome sequence and tools for interrogating individual genomes provide an unprecedented opportunity to apply genetics to medicine. Mendelian conditions, which are caused by dysfunction of a single gene, offer powerful examples that illustrate how genetics can provide insights into disease. Cystic fibrosis, one of the more common lethal autosomal recessive Mendelian disorders, is presented here as an example. Recent progress in elucidating disease mechanism and causes of phenotypic variation, as well as in the development of treatments, demonstrates that genetics continues to play an important part in cystic fibrosis research 25 years after the discovery of the disease-causing gene.

Cystic fibrosis (OMIM 219700) is a life-limiting autosomal recessive disorder that affects 70,000 individuals worldwide. The condition affects primarily those of European descent, although cystic fibrosis has been reported in all races and ethnicities. Abnormally viscous secretions in the airways of the lungs and in the ducts of the pancreas in individuals with cystic fibrosis cause obstructions that lead to inflammation, tissue damage and destruction of both organ systems (FIG 1). Other organ systems containing epithelia -such as the sweat gland, biliary duct of the liver, the male reproductive tract and the intestine -are also affected. Loss of pancreatic exocrine function results in malnutrition and poor growth, which leads to death in the first decade of life for most untreated individuals. Replacement of pancreatic enzymes and intensive therapy guided by multidisciplinary teams have revolutionized the treatment of cystic fibrosis, resulting in progressive improvements in survival to a median predicted age of 37 years for children born with cystic fibrosis today¹. Obstructive lung disease is currently the primary cause of morbidity and is responsible for 80% of mortality².

Twenty-five years ago, a variant (p.Phe508del; also known as F508del in legacy nomenclature) in the cystic fibrosis transmembrane conductance regulator (*CFTR*) gene was found to be the most common cause of cystic fibrosis³⁻⁵. Demonstration that *CFTR* functions as a chloride channel regulated by cyclic AMP (cAMP)-dependent phosphorylation⁶ was consistent with the ion transport disturbances documented in cystic fibrosis tissues. Key insights into cystic fibrosis pathophysiology were derived from the

study of CFTR mutants⁹, correlation of CFTR dysfunction with the cellular manifestations of cystic fibrosis¹⁰ and elucidation of protein partners involved in biogenesis and membrane function. Identification of disease-causing variants in *CFTR* contributed a tool for both the diagnosis of cystic fibrosis and the identification of cystic fibrosis carriers¹², demonstrated the degree to which CFTR dysfunction correlates with clinical features¹ and revealed that CFTR dysfunction can create phenotypes other than cystic fibrosis¹⁴. Over the past 5 years, there has been remarkable progress in the development of small-molecule therapy targeting CFTR bearing select disease-causing variants^{15,16}.

The purpose of this Review is to highlight advances over the past decade in our understanding and treatment of cystic fibrosis that were informed by genetics. Given the breadth of the cystic fibrosis field, not all of the important contributions and publications relevant to the topic can be included. Examples have been chosen to illustrate that genetics continues to have a role in the research of Mendelian disorders long after the causative variants and the responsible gene have been discovered. This Review covers new insights into the processing defect caused by the F508del variant, advances in stem cell technology that can enable testing of therapeutics for a wide range of *CFTR* genotypes and the development of new animal models that are informing our understanding of organ pathology in cystic fibrosis. I also summarize progress in parsing genetic and nongenetic contributions to variability in cystic fibrosis and in the identification of modifier loci. The final section describes efforts to determine the molecular and phenotypic consequences of the majority of cystic fibrosis-causing variants and to develop molecular treatments for every defect in *CFTR*.

Insights into disease mechanism

Molecular basis of CFTR dysfunction. Almost 2,000 variants have been reported to the Cystic Fibrosis Mutation Database, one of the first and most successful locus-specific databases. Among these variants, 40% are predicted to cause substitution of a single amino acid, 36% are expected to alter RNA processing (including nonsense, frameshift and missplicing variants), 3% involve large rearrangements of *CFTR*, and 1% affects promoter regions; 14% seem to be neutral variants, and the effect of the remaining 6% is unclear. Disease-causing variants can affect the quantity and/or function of CFTR at the cell membrane (FIG 2). Historically, *CFTR* variants have been grouped into five (and sometimes six) functional classes⁹. The class system provides a useful framework for understanding the primary defect at the cellular level. However, binning of variants into one class is problematic, as multiple processes can be affected by a single variant. For example, F508del causes aberrant folding of CFTR and subsequent degradation of the majority of the synthesized protein¹⁷. The minor fraction of F508del-CFTR that is trafficked to the cell membrane has severely reduced membrane residency and aberrant chloride channel function¹⁸. Furthermore, the three-nucleotide deletion responsible for the F508del variant also causes a synonymous change in the triplet that encodes isoleucine at codon 507 (ATC-7ATT). The change alters the structure of the F508del-CFTR mRNA, which leads to a reduction in translation efficiency¹⁹. Thus, F508del could be assigned to at least three classes. Various missense variants also cause defective processing and alter chloride channel

function of CFTR^{20,21}. Appreciating the diversity of effects caused by a *CFTR* variant is important in the design of molecular treatments for cystic fibrosis (see below).

Disease-causing variants provide a reservoir of naturally occurring deleterious amino acid deletions and substitutions that have proved to be informative for dissecting the tertiary structure of CFTR. CFTR is composed of three major motifs: domains that interact with ATP, termed nucleotide-binding domain 1 (NBD1) and NBD2; regions that anchor the protein in the membrane known as membrane-spanning domain 1 (MSD 1) and MSD2; and an area containing numerous sites for phosphorylation called the regulatory domain (also known as the R domain) (FIG 2). It had been recognized for some time that deletion of phenylalanine at codon 508 (F508) causes instability of NBD 1, but how this localized structural defect causes misfolding of the entire protein was poorly understood²². Furthermore, it was known that disease-causing missense variants outside the NBD 1 domain - notably, a cluster in the fourth cytosolic loop (CL4) within MSD2 -also cause misfolding of the protein^{22,4}. Modelling based on the atomic structure of related proteins^{25,26} and cysteinecrosslinking experiments²⁷ revealed an interaction between the NBDs and MSDs of CFTR. Notably, F508 occurs at an interface between NBD1 and CL4, and seems to be capable of forming hydrogen bonds with arginine at codon 1070 (R1070)²⁶. Restoration of NBD 1 assembly using suppressor mutations produces only partial recovery of CFTR processing, which indicates that the F508del variant also affects interactions elsewhere in the full-length protein²⁸. Intriguingly, introduction of the disease-associated p.Arg1070Trp (legacy R1070W)²⁹ variant in CL4 and correction of NBD 1 misfolding using synthetic suppressor mutations could restore processing to F508del-CFTR^{28,30,31}. These findings lay the groundwork for structure-based selection of molecules that correct the processing defects in CFTR caused by disease-causing variants³².

Tissue culture. Assessment of the functional consequence of variants requires the appropriate cellular context. For CFTR, primary epithelial cells are the most relevant system; however, these cells are short-lived and require accessions from internal organs such as the lung . Furthermore, availability and compliance substantially limit acquisition of primary cells from individuals with rare *CFTR* genotypes. Recent advances in stem cell biology have provided new methods for generating well-differentiated cell lines from individuals with cystic fibrosis. Human embryonic stem cells and intestinal stem cells from individuals with cystic fibrosis have been coaxed into differentiating into secretory epithelial cells that manifest defects in CFTR-mediated chloride transport^{34,35}. Small-molecule correctors for the F508del variant were efficacious in restoring CFTR function in both cell types, which showed the utility of these systems for evaluating therapeutics^{32,34}. Genetic reversion of the F508del variants to wild type using CRISPR-Cas9 editing has been achieved in intestinal organoids³⁶. Conversion of well-differentiated epithelial cells from human ectocervix and trachea into 'conditionally reprogrammed' cells using Rho kinase inhibitor offers an alternative approach to derive individual-specific cell types³⁷. These methods provide new tissue models for examining function and dysfunction of CFTR from individuals with specific *CFTR* genotypes.

Animal models. One of the most useful tools in the investigation of genetic disorders is animal models. Cystic fibrosis is unique among human genetic disorders in that five animal

models (mouse³⁸, rat³⁹, ferret⁴⁰, pig⁴¹ and zebrafish⁴²) have been created. Although these animal models have not replicated human cystic fibrosis precisely, their differences have proved to be highly instructive for understanding disease mechanisms at the organ-system level (TABLE 1). The *Cfr* gene in mice has been extensively manipulated to derive lines that do not express CFTR and lines that express CFTR bearing variants equivalent to those observed in humans (for example, F508del and p.Gly551Asp (legacy G551D))⁴. Even though airway epithelial cells display ion transport abnormalities that are consistent with loss of CFTR function, overt lung disease is not evident in newborn or young mice with cystic fibrosis⁴⁴. The absence of lung disease similar to that seen in humans with cystic fibrosis has been ascribed to the presence of alternative pathways for chloride transport in mouse epithelial cells⁴⁵. This observation suggests that ion channels other than CFTR might be exploited to recover chloride transport in cystic fibrosis cells. Indeed, a suitable candidate may have already been identified in *TMEM16A* (also known as *ANO1*), which encodes a calcium-activated chloride channel first identified in mouse airways⁴⁶. Despite the phenotypic differences between mice and humans, the study of cystic fibrosis mouse models has provided invaluable insights into the role of other ion channels in the development of lung disease, the biology of intestinal obstruction and the evaluation of candidate modifiers⁴⁷, and such models also provide an *in vivo* platform for testing therapeutics⁴³.

To create animal models that are more likely to recapitulate disease mechanisms operating in the lung, investigators have selected species on the basis of specific anatomical features. Of particular importance are the number and distribution of airway submucosal glands — a site of high CFTR expression - in humans⁴⁸. The lungs of pigs and ferrets represent reasonable approximations of human lung architecture, and successful *CFTR* knockouts have been achieved in both species using homologous recombination^{40,49}. The porcine model of cystic fibrosis develops lung disease comparable to that observed in humans, albeit at an earlier stage of life⁵⁰. Consequently, the cystic fibrosis pig provides an opportunity to address the sequence of events in early-stage lung disease and to evaluate therapeutics in new-born animals. One of the key issues in the early stages of human cystic fibrosis is the genesis of inflammation in the lungs. Some studies suggest that infection of the airways trigger inflammation, whereas others demonstrate the presence of an inflammatory response in the absence of lung infection⁵¹. This distinction is important, as treatment strategy of each scenario is markedly different. In the lungs of cystic fibrosis pigs, inflammatory markers are not elevated until polymicrobial infection ensues⁵⁰. Furthermore, pigs with cystic fibrosis exhibit an inability to eradicate bacterial pathogens, which may be partly due to abnormalities in the pH of the airway surface liquid caused by reduced CFTR-dependent bicarbonate transport⁵². Reduction in the size of the large airways in newborn pigs with cystic fibrosis has been associated with air trapping, a defect noted in human infants with cystic fibrosis⁵³. Congenital anomalies in tracheal rings in mice, pigs and humans with cystic fibrosis (TABLE 1) support the concept that defects in the development of the pulmonary tree contribute to early-stage lung disease in cystic fibrosis. Finally, ion transport studies in newborn cystic fibrosis pigs have questioned the longstanding concept that sodium absorption is increased in the airway epithelia⁵⁴. The cystic fibrosis pig model provides compelling evidence that loss of chloride and bicarbonate transport,

maldevelopment of the airways and infection are the important drivers of early-stage lung disease in cystic fibrosis.

Box 1

Twin studies for estimating heritability

A twin study represents a naturally balanced, age-matched design that provides a powerful method for determining the contribution of genetics to a trait. This approach capitalizes on the different roles of variant sharing among monozygotic twins (who are presumed to be 100% identical except for *de novo* mutations), and dizygotic twins and siblings (who share ~50% of their variants). Environmental exposures are similar (but not identical) *in utero* and continue to be similar as twins grow in the same household. Thus, by comparing the degree of similarity among monozygotic twin pairs to dizygotic twin pairs, an estimate can be obtained of the degree to which trait variance can be attributed to genetic variation (that is, heritability)¹¹⁷. Heritability estimates generated in this manner range from 0 to 1. Estimates approaching 1.0 indicate strong genetic influence, whereas those near zero exclude a prominent role for genetic variation. The effect of shared environment can be estimated by comparing trait variance among twin pairs living together to those living apart. Finally, within-pair variance for monozygotic twins provides a measure of unique environmental and stochastic components, as the genetic variance is, by definition, almost zero between monozygotic twins. As the number of twin pairs available for study of a Mendelian disorder is generally small, only crude estimates of genetic control of trait variance can be obtained. Nevertheless, the approach outlined above is widely applicable, as family-based studies can be used to estimate genetic control of traits that constitute any Mendelian disorder.

Mammalian models of cystic fibrosis have also provided insights into disease processes in other affected organ systems. Pancreatic exocrine dysfunction is closely correlated with the development of neonatal intestinal obstruction in humans with cystic fibrosis. However, animal models exhibit minimal (mouse) to severe (pig) pancreatic exocrine disease compared with humans, yet the incidence of intestinal obstruction is higher in all mammalian models than in humans (TABLE 1). Recovery of intestinal expression of CFTR prevents obstruction in cystic fibrosis mice, pigs and ferrets^{40,55,56}. These two observations suggest that loss of CFTR function in the intestine rather than pancreatic exocrine dysfunction is the primary cause of intestinal obstruction in cystic fibrosis. Diabetes mellitus is an age-dependent complication that affects 40% of individuals with cystic fibrosis by 35 years of age²; this disorder is also closely correlated with pancreatic exocrine dysfunction. Destruction of the exocrine pancreas has been proposed to stress the endocrine pancreas, which leads to loss of insulin-secreting cells⁵⁷. However, features of diabetes occur before the development of severe pancreatic exocrine disease in ferrets⁵⁸ and in the absence of substantial loss of insulin-producing cells in cystic fibrosis pigs⁵⁹. Both observations suggest that cystic fibrosis-related diabetes is the result of an intrinsic defect in the endocrine pancreas caused by loss of CFTR function.

Variation in disease severity

The relative contribution of genetic modifiers. Individuals with cystic fibrosis show a high degree of variability in disease severity, complications and survival. It was initially postulated that a substantial fraction of phenotypic variability would be explained by allelic heterogeneity in the dysfunctional gene⁶⁰. *CFTR* genotype correlates well with pancreatic exocrine disease severity and modestly with sweat chloride concentration^{61,62}. However, it has been difficult to detect a relationship between lung function and *CFTR* genotype^{61,63}, with a few notable exceptions⁶⁴. Analyses of families with affected twins (BOX 1) have quantified the degree to which variables beyond *CFTR* — such as genetic modifiers, environmental factors and/or stochasticity -influence variability in lung disease severity (FIG 1). Affected monozygotic twin pairs exhibit greater similarity for lung function than affected dizygotic twin and sibling pairs (siblings were used as a proxy for dizygotic twins)^{65,66}. By comparing clinical measures of affected twin pairs when they lived together to the same measures after they moved apart, 50% of the difference in lung function measures could be attributed to genetic modifiers. The remaining variation was due to environmental exposures, primarily those unique to each individual, and to stochastic factors⁶⁷. Together, these family-based studies demonstrate that genetic modifiers have considerable influence on lung function variation in cystic fibrosis.

The contribution of genetic modifiers to four other traits that are relevant to survival in cystic fibrosis has been estimated (FIG 1). Chronic colonization of the lungs with the bacterial pathogen *Pseudomonas aeruginosa* is a feature of advancing lung disease in cystic fibrosis and is associated with reduced survival⁶⁸. Establishment of chronic *P aeruginosa* infection and age at establishment are highly influenced by genetic factors⁶⁹. Poor growth is a hallmark of cystic fibrosis owing to pancreatic exocrine disease and deficiency of insulin-like growth factor 1. Although replacement of pancreatic digestive enzymes has improved nutritional status of individuals with cystic fibrosis, those with extremely low body mass index (BMI) remain challenging to treat. Genetic control of BMI independent of *CFTR* seems to be substantial, as estimated heritability ranges from 0.54 to 0.8 (REFS 70,71). Affected-twin analysis also revealed that genetic modifiers are primarily responsible for the age at onset of diabetes (heritability: 1.0; confidence interval: 0.42-1.0). Diabetes mellitus is associated with more rapid decline in lung function in individuals with cystic fibrosis, and medical management of glucose levels improves survival¹⁷. Finally, obstruction of the small intestine (a condition known as meconium ileus) complicates the management of 15% of newborns with cystic fibrosis. Strain-specific differences in the rate of intestinal obstruction in cystic fibrosis mice first showed that this trait could be modified by genes other than *Cftr*⁷⁴. Analysis of twins with cystic fibrosis demonstrated that genetic modifiers have the predominant role in the development of intestinal obstruction (heritability: 1.0)⁷⁵.

Finding variants that modify cystic fibrosis. Extensive understanding of cystic fibrosis pathophysiology presents an opportunity to interrogate candidate genes as potential modifiers. In the lungs, loss of CFTR leads to exuberant inflammation, neutrophil recruitment, tissue damage and eventual replacement with fibrotic connective tissue. At least 50 genes encoding proteins that participate in these cellular and tissue functions have been investigated as candidate modifiers of lung disease severity in cystic fibrosis⁷⁶. As the

results of candidate modifier gene studies have been extensively reviewed elsewhere^{76,80}, this Review highlights the insights gained from the study of one candidate gene that illustrates the potential and challenges in dissecting the complex interactions that modify disease severity.

A key element of lung disease in cystic fibrosis is the response to recurrent tissue injury. Transforming growth factor beta 1 (*TGFBI*) was an intriguing candidate for a modifier of cystic fibrosis lung disease, as it is involved in tissue repair and extracellular matrix production. Furthermore, variants in *TGFBI* have been associated with risk of asthma and chronic obstructive pulmonary disease (COPD) -two conditions that have features in common with cystic fibrosis lung disease. Two variants that increase levels of *TGFBI* were correlated with severe lung disease as determined by airway flow measurements⁸¹. However, the direction of effect differs between cystic fibrosis and COPD; the same alleles show a deleterious consequence in one disorder but a protective effect in the other. It has been speculated that the presence of functional *CFTR* may invert the clinical consequences of increased *TGFBI* expression in individuals with COPD⁸¹. Gene-gene and gene-environment studies using *TGFBI* variants have begun to address the issue of context. An obvious first place to look for gene-gene interaction is between *TGFBI* modifier variants and *CFTR* disease-causing variants. Drumm, Knowles and colleagues established *TGFBI* as a modifier primarily in individuals who are homozygous for the common cystic fibrosis-causing variant F508del⁸¹. A subsequent study demonstrated that the alternative alleles of the same *TGFBI* variants were associated with less severe lung disease, but the effect was limited to individuals with *CFTR* genotypes other than F508del homozygosity⁸². Interaction has been reported between *TGFBI* and mannose-binding lectin 2 (*MBL2*), which is another genetic modifier of cystic fibrosis. Variants associated with increased *TGFBI* expression amplify the deleterious effects of *MEL* deficiency upon lung infection and airway deterioration⁸⁴. Environmental context is also important, as variants in *TGFBI* exacerbate the pernicious effects of exposure to second-hand smoke in patients with cystic fibrosis⁸⁵. These observations indicate that deducing the clinical effect of a modifier variant greatly depends on the context in which it occurs.

To identify novel modifiers, research groups have combined patient populations to achieve sufficient power in genome-wide methods. Formation of a North American Cystic Fibrosis Gene Modifier Consortium — composed of investigators at the University of North Carolina, USA; the Hospital for Sick Kids in Toronto, Canada; and Johns Hopkins University in Baltimore, Maryland, USA - facilitated both association and linkage studies on 3,500 individuals. Although the three sites used different study designs, each agreed to use the same measure of lung disease severity, thereby enabling an analysis of unrelated subjects recruited by the centres in North Carolina and Toronto, as well as replication in related subjects participating in the Johns Hopkins study. In a genome-wide association study (GWAS), a significant region was detected between two genes on chromosome 11: *EHF*, which encodes an epithelial transcription factor, and *APIP*, which encodes an inhibitor of apoptosis⁸⁶. The known functions of *EHF* and *APIP* suggest biologically plausible roles in modifying lung function in cystic fibrosis⁸⁶. To search for rare variants that modify lung function, linkage analysis was carried out on 486 affected sibling pairs. A locus on

chromosome 20q13.2 harbouring only 4 genes was identified⁸⁶. The identification of each locus demonstrates the feasibility of genome-wide approaches for uncovering new pathways that modify disease severity in cystic fibrosis.

Searching for genetic modifiers of other cystic fibrosis traits has provided mechanistic insights on several fronts. First, risk variants for other diseases operate as modifiers of similar conditions in patients with cystic fibrosis. Variants in four genes that confer risk for type 2 diabetes mellitus on the general population (*TCF7L2*, *CDKALI*, *CDKN2A/B* and *IGF2BP2*) modify age at onset of diabetes in cystic fibrosis⁸⁷(FIG 1). The Z allele of *SERYINA*, which causes alpha-1 antitrypsin deficiency and confers risk for emphysema and liver disease, modifies risk of cirrhotic liver disease in cystic fibrosis⁸⁸. Second, modifiers exhibit pleiotropic effects on the cystic fibrosis phenotype. Variants in *SLC26A9* - which encodes a chloride and bicarbonate channel that interacts with CFTR - modify risk for neonatal intestinal obstruction and diabetes⁸⁹. Furthermore, solute carriers associated with risk of neonatal intestinal obstruction also modify lung disease severity in young patients with cystic fibrosis (*SLC9A3* and *SLC6A14*) and age at first infection with *P. aeruginosa* (*SLC6A14*)⁹⁰. Third, informing GWASs with knowledge of CFTR function increases the yield of significant associations. This approach, termed hypothesis-driven GWASs, revealed that proteins residing in the same cellular location as CFTR are enriched for modifiers of neonatal intestinal obstruction. Fourth, exome sequencing can find rare modifier variants. Variants in *DCTN4* - which encodes a dynactin protein involved in autophagy- are associated with age at onset of chronic infection with *P. aeruginosa*⁹¹. The identification of *DCTN4* as a modifier may provide a mechanistic link to defective autophagy in cystic fibrosis cells⁹². Conversely, loss-of-function variants in *CFTR* have been linked to a variety of other conditions (BOX 2). Together, these findings illustrate that the complex mechanisms underlying trait modification in cystic fibrosis can be informed by genetics, especially if approaches beyond standard association and linkage are undertaken.

Molecular diagnosis and therapy

Screening, diagnosis and functional annotation of CFTR variants. For almost 2 decades, a panel of 23 of the most common variants - vetted by an expert committee of the American College of Medical Genetics — has been used to diagnose cystic fibrosis and to screen for carriers and affected newborns. Expansion of the panel to increase sensitivity remained challenging, and the disease liability for most of the remaining *CFTR* variants was not known⁹⁴. The Clinical and Functional Translation of *CFTR* (CFTR2) project was initiated in 2010 to increase the number of annotated *CFTR* variants. Clinical and *CFTR* genotype data collected on 39,696 individuals enrolled in cystic fibrosis patient registries and clinics in North America and Europe were used to determine the penetrance of variants for cystic fibrosis. To widely and rapidly disseminate results, features associated with each variant are available on a public website (see CFTR2). Content is tailored to educate patients, family and the public about the clinical implications that can and cannot be inferred from *CFTR* genotype. Inclusion of 127 *CFTR* variants annotated as disease-causing in screening assays increased the sensitivity for detection of cystic fibrosis alleles in white European individuals from 85% to 95%²⁰. Although this improvement seems modest, the recessive nature of cystic fibrosis requires ascertaining variant status in two *CFTR* genes. Thus, at

95% sensitivity, only 0.25% of individuals with cystic fibrosis will have neither disease-causing variant identified. Translation of these findings should improve the accuracy of screening programmes, and will aid diagnosis and treatment, as 99.75% of individuals with cystic fibrosis should carry at least one disease-causing *CFTR* variant. Owing to population differences in the frequency of *CFTR* variants, in particular the F508del variant, the sensitivity of screening is lower in non-white individuals. Inclusion of affected individuals from South America, Africa, the Middle East and East Asia in the current phase of *CFTR2* recruitment will provide a more complete inventory of variants found in non-white populations. As of late 2014, *CFTR2* has obtained data on 73,000 individuals, thereby exceeding the estimated number of individuals with cystic fibrosis worldwide (70,000). Given the completeness of the ascertainment, the sensitivity of screening in all populations will be improved, substantially in some cases, once variant annotation is completed.

Molecular therapy for cystic fibrosis. The most exciting development in cystic fibrosis research since the identification of *CFTR* has been the successful implementation of therapy that augments the function of mutant *CFTR*. Increasing the chloride channel activity of mutant forms of *CFTR* was shown to be a viable therapeutic approach two decades ago⁹⁵. Development of compounds that selectively activated *CFTR* at doses that could be achieved *in vivo* proved to be difficult, and attention was focused on gene replacement methods for the next decade. However, the search for a molecular correction of *CFTR* has not been abandoned. A unique partnership between a biotech company and the US Cystic Fibrosis Foundation initiated empirical screens for small molecules that target *CFTR* mutants. A promising compound termed ivacaftor (also known as VX-770 and Kalydeco (Vertex Pharmaceuticals)) increased chloride transport of primary airway cells bearing the G551D variant up to 50% of wild-type level⁹⁶. Significant but more modest increases in G551D-*CFTR* activity were observed in immortalized cell lines (10-30% of wild-type *CFTR*)^{97,98}. The increase in *CFTR* function observed in cell lines achieved levels that would be expected to produce a clinical response in lung function (BOX 3). Indeed, Phase III clinical trials over 4-week and 48-week intervals demonstrated that ivacaftor improved lung function (by 10% on average) and reduced sweat chloride concentration (to an average concentration below the diagnostic threshold of 60 mM) in individuals with cystic fibrosis carrying the G551D variant^{15,99} (FIG 3a). Assessment of *in vivo* response to ivacaftor using sweat volume measures in 5 patients estimated recovery of 1.6-7.7% of wild-type *CFTR* function, while estimates derived from sweat chloride concentration and nasal potential difference measurements in 39 individuals indicated recovery of 35-40% of normal *CFTR* function^{100,101}.

Box 2

CFTR in other diseases

Male infertility

The concept that dysfunction of cystic fibrosis transmembrane conductance regulator (*CFTR*) could create disorders other than cystic fibrosis was first illustrated by obstructive male infertility due to congenital bilateral absence of the vas deferens (CBAVD; OMIM 277180)¹¹⁸. The anatomical features of CBAVD are identical to those seen in moles with

cystic fibrosis, and some individuals with CBAVD have subtle features of cystic fibrosis, such as mildly elevated sweat chloride concentration or minimal airway disease¹¹⁹. However, a fraction of moles with CBAVD manifest no evidence of cystic fibrosis in detailed studies of the lungs, pancreas and sinonasal gland¹²⁰. Furthermore, the distribution of *CFTR* variants differs between CBAVD and cystic fibrosis, and a much higher fraction of variants is associated with residual function occurring in CBAVD¹²¹. Thus, CBAVD is part of the cystic fibrosis spectrum caused by *CFTR* dysfunction (FIG. 1), but it is also viewed as clinically distinct, particularly in moles with features limited to the vas deferens.

Pancreatitis

Loss-of-function variants in *CFTR* have also been linked to a variety of conditions collectively termed 'CFTR-related disorders', as reviewed by others^{14,122}. Discovery of a pathological role for *CFTR* has proved to be particularly instructive for the study of pancreatitis. Pancreatitis is a known complication of cystic fibrosis, primarily occurring in individuals with preserved pancreatic exocrine function¹²³. Pancreatitis in the general population is a heterogeneous disorder with heritable and idiopathic sporadic forms¹²⁴. A subset of heritable forms of recurrent acute and chronic pancreatitis can be attributed to *CFTR* dysfunction¹²⁴. As with CBAVD, the distribution of *CFTR* variants differs from that of cystic fibrosis. Recent evidence suggests that cells expressing *CFTR* bearing variants associated with pancreatitis, but not with cystic fibrosis, manifest defective bicarbonate transport, while chloride channel function is preserved¹²⁵. This concept aligns well with the role of *CFTR* as an important mediator of bicarbonate transport in the pancreatic ducts¹²⁶.

Manifestations in cystic fibrosis carriers

CFTR variants also act as risk alleles for multigenic disorders in the general population. Idiopathic disseminated bronchiectasis is a relatively rare pulmonary airway disease that manifests features similar to those observed in the lungs of individuals with cystic fibrosis. Several studies have shown a higher frequency of deleterious *CFTR* variants in individuals with bronchiectasis than in control subjects¹²⁷. Bronchiectasis is complicated by infection with non-tuberculous mycobacteria or with the fungus *Aspergillus fumigatus*, which has also been associated with an increased frequency of *CFTR* variants^{127,128}. Chronic rhinosinusitis (CRS) is an aetiologically heterogeneous condition affecting ~15% of the general population in the United States. CRS is a common complication in individuals with cystic fibrosis. Genotyping of subjects meeting rigorous criteria for CRS revealed an excess of carriers of a single deleterious *CFTR* variant compared to disease-free controls¹²⁹. In these studies, the entire coding region of *CFTR* was examined to exclude a second deleterious variant. Support for the concept that presence of a single loss-of-function variant in *CFTR* predisposes to CRS was derived from the observation that the obligate heterozygous carriers of the deleterious *CFTR* variant (that is, the parents of individuals with cystic fibrosis) had a threefold increase in prevalence of CRS¹³⁰.

Assessing whether pancreatitis, bronchiectasis or sinusitis can be attributed to *CFTR* dysfunction in a heterozygous cystic fibrosis carrier requires detailed phenotyping to

exclude other conditions, including mild forms of cystic fibrosis¹³¹. This challenge will become particularly pertinent as we enter into an age where CFTR dysfunction can be treated at the molecular level. Furthermore, associating *CFTR* variants with common multigenic disorders has substantial implications, as there are an estimated 20 million heterozygous carriers of cystic fibrosis in the world. Many carriers of deleterious *CFTR* variants are becoming aware of their status as population testing is widespread in the United States and is becoming more common in Europe. However, the penetrance of most *CFTR* variants for the traits discussed above is not known. Establishing the penetrance of variants in disease-associated genes is a considerable and important challenge¹³²

Box 3

Reversing cystic fibrosis: how much CFTR is needed?

Effective treatment of cystic fibrosis at the molecular level requires restoration of cystic fibrosis transmembrane conductance regulator (CFTR) function in affected tissues. Genotype and phenotype correlation revealed that organ systems have different requirements for CFTR function¹³. Splice site variants that reduce but do not eliminate production of some CFTR mRNA have been particularly informative in this regard¹³³. Improvement in lung function and other features of cystic fibrosis (for example, SiNE chloride concentration) occurs at CFTR mRNA levels above 5% of normal^{134,135}. Splice site variants that allow CFTR transcript levels to reach 10-25% of normal levels have been found in individuals that do not have cystic fibrosis lung disease^{136,137}. As the residual RNA transcript is of full length, it has been assumed that the quantity of transcribed CFTR protein will correspond to the level of remaining full-length CFTR transcripts. A lower boundary of 10% of normal levels for relieving pulmonary disease is supported by cell mixing experiments, which indicate that airway epithelial ion transport is normalized when 6-10% of cells have corrected CFTR function¹³⁸. Rescue of other key functions of the respiratory epithelium may require higher levels of CFTR function. Restoration of mucus transport required ~25% of cells to be corrected¹³⁹. These estimates represent averages; variation in genetic and non-genetic modifiers is likely to broaden the range of CFTR correction required at the individual level.

The successful clinical deployment of ivacaftor encouraged the search for compounds that can correct other defects in CFTR. The prevalence of the F508del variant has attracted intense interest in reversing the folding defect caused by this variant. Screening of small molecules followed by chemical modification of active compounds led to the formulation of lumacaftor (also known as VX-809), which increased chloride transport of primary airway cells bearing F508del-CFTR to 14% of wild-type levels¹⁰². A clinical trial of lumacaftor produced a dose-dependent improvement in CFTR function measured in the sweat gland of patients carrying the F508del variant¹. However, CFTR function was not augmented in nasal epithelia, and lung function measures were not improved¹ (FIG 3a). As ivacaftor confers wild-type levels of open probability on F508del-CFTR, albeit as a result of a different pattern of channel gating⁹⁶, clinical trials combining ivacaftor with the corrector

lumacaftor have been undertaken. This approach follows the reasonable logic that a potentiator can increase the activation of 'corrected' F508del-CFTR, thus amplifying the effect achieved by each compound alone¹⁰² (FIG 3b). A Phase II clinical trial demonstrated that the combination of lumacaftor and ivacaftor improved measures of lung function and sweat chloride concentration in individuals homozygous for the F508del variant¹⁶. This encouraging result has been followed by the announcement that 2 Phase III clinical trials of combined lumacaftor and ivacaftor involving 1,100 F508del homozygotes over a 24-week period documented improvement in lung function (See Vertex press release). Although the changes in lung function measures were modest, there were concurrent improvements in secondary end points, providing encouraging evidence of clinical efficacy. However, two groups have recently reported that ivacaftor exposure for 48 hours diminishes the correction of F508del-CFTR conferred by lumacaftor in primary and immortalized cells^{104,105}. Reduction in the quantity of 'corrected' F508del-CFTR due to ivacaftor may explain the modest responses observed in the clinical trials (FIG 3b). Thus, compounds selected for combinatorial therapy will have to be carefully screened for undesirable interactions. With a panel of 'correctors' and 'potentiators' in hand, screening can proceed for combinations that act cooperatively^{32,105} (FIG 3b). Furthermore, the available drugs could be used to screen for novel compounds that interact synergistically to further recover function of F508del-CFTR¹⁰⁶ (FIG 3b).

Although F508del and G551D account for a large fraction of cystic fibrosis alleles, 7% of patients with cystic fibrosis carry neither variant. To extend available therapy to as many individuals as possible, variants that permit translation of the CFTR protein can be evaluated for response to clinically approved corrector and potentiator compounds. However, as hundreds of translatable variants have been found in *CFTR*, it will be challenging to perform clinical efficacy studies of all variants, and many of these variants are carried by only a few affected individuals. Therefore, a new approach is required to extend approved efficacious treatments to individuals with rare variants in *CFTR*. It has been proposed to group *CFTR* variants into theratypes according to their effect on the CFTR protein and in response to corrector and potentiator compounds. Previously unclassified variants can be provisionally assigned to theratypes on the basis of their effect on CFTR quantity and function in cell-based studies. Response of CFTR bearing the unclassified variant to the profile of compounds that define the theratype would confirm that the assignment is appropriate (FIG 3c). Measures of *in vivo* CFTR function can then be used to verify clinical response and to justify ongoing treatment of individual patients¹⁰⁷. To this end, 9 missense variants were shown to cause a defect in activation (that is, gating) of CFTR and reduction in chloride transport ranging from 0% to 9.7% of normal levels¹⁰⁸. As noted for the G551D variant, ivacaftor treatment of immortalized cells expressing CFTR bearing each of these variants increased chloride transport from 21% to 157% of normal levels¹⁰⁸, thereby predicting that clinical response should occur in individuals carrying these variants (FIG 3c). Subsequently, a Phase III clinical trial of 39 individuals demonstrated that ivacaftor was efficacious for 8 of the 9 variants, leading to rapid approval by the US Food and Drug Administration (NDA 203188). The four individuals carrying one variant did not respond sufficiently to warrant approval for ivacaftor treatment. The rapid expansion of small-molecule therapy for cystic fibrosis, from cell-based studies to clinical application, provides a new paradigm for drug

development for genetic disorders and an excellent example of the promise of personalized medicine.

To treat all patients with cystic fibrosis, it will be necessary to address variants that prevent or severely decrease production of the CFTR protein through alterations in RNA processing. Nonsense and frameshift variants that introduce a premature termination codon (PTC) pose several hurdles that must be cleared to achieve therapeutic quantities of the protein. Most PTC variants invoke RNA degradation through the nonsense-mediated RNA decay (NMD) pathway¹⁰⁹. Counteracting NMD to stabilize mRNA *in vivo* is challenging owing to the possibility of off-target effects. Synthesis of a full-length protein requires readthrough of the PTC, and some success in suppressing nonsense variants in *CFTR* using compounds derived from aminoglycosides has been achieved¹¹¹ (FIG 3d). Clinical trials of PTC suppressors have documented modest recovery of CFTR function in the nasal epithelia, although improvement in lung function has not been reported¹¹². Even when readthrough is successful, the incorporation of a non-native amino acid at the location of the nonsense variant could affect protein processing and function¹⁰⁹. Thus, augmenting the function of CFTR with ivacaftor following PTC suppression seems to be a viable approach to exceed the therapeutic threshold (FIG 3d). Alternatively, strategies such as induction of the unfolded protein response could be used to attenuate NMD, thereby stabilizing transcripts with PTCs for possible readthrough¹¹⁴. Variants that cause aberrant RNA splicing pose a different set of challenges. On the one hand, suppression of variants that activate cryptic splice sites, such as c.3717 + 12191 C→T (legacy 3849 + 10kbC-7T), can substantially increase the amount of normally spliced RNA transcript and protein (FIG 3d). On the other hand, variants that alter canonical nucleotides in splice sites are proving difficult to treat, although manipulation of splicing factors (for example, U1 small nuclear RNA) and the splicing process (for example, *trans-splicing*) shows some promise¹¹⁵. The remaining variants are rearrangements, which require replacement of one or more exons or the entire coding sequence of *CFTR*. Transfer of DNA to epithelial cells has been extensively explored as a therapeutic approach for cystic fibrosis, but efficient delivery to airway cells *in vivo* remains problematic¹¹⁶. Finally, although molecular treatment is at hand, variation in response^{15,35} indicates that underlying individual differences will have to be addressed to achieve the goal of attaining a normal lifespan for all patients with cystic fibrosis.

Conclusions

The discovery of *CFTR* 25 years ago was a triumph for genetics and a potent demonstration of its ability to deliver the molecular culprit in a Mendelian disorder. Cystic fibrosis is now positioned to reap the dividends of personalized medicine as variant-specific therapy is deployed, and a growing understanding of the genetic and environmental modifiers of cystic fibrosis enables targeting of individual risk factors. The development of new genetic models of cystic fibrosis in pigs, ferrets, rats and zebrafish provides opportunities to investigate pathophysiology and to explore therapies at the earliest stages of disease. Newborn and population screening enables prospective management of affected individuals from birth, and genomic variation will provide information on the trajectories that individual patients are likely to follow. Genetics has played and will continue to play a key part in achieving a normal lifespan for individuals with cystic fibrosis.

Acknowledgements

The author thanks P. Thomas, D. Sheppard, M. Knowles, M. Drumm and B. Guggino for providing commentary and critique of this Review, and members of the CFTR2 team (C. Penland, J. Rommens, C. Castellani and M. Corey) for many insights regarding the clinical and functional consequences of CFTR variants. He also thanks P. Durie, H. Corvol and the members of the International Cystic Fibrosis Modifier Consortium for discussions about modifiers of cystic fibrosis, and members of the Cutting laboratory, especially P. Sosnay, S. Blackman, J. M. Collaco and K. Raraigh, for contributions to concepts presented in this Review. The author's work is supported by grants 5R01DK044003 from the NIDDK, and grants CUTTING08A, CUTTING09A and CUTTING10A from the US Cystic Fibrosis Foundation. Competing interests statement The authors declare competing interests: see Web version for details.

Glossary

Pancreatic exocrine	Pertaining to the portion of the pancreas that produces digestive enzymes that are combined with alkaline secretions from the pancreatic ducts and secreted into the intestine to aid digestion.
Locus-specific databases	Collections of DNA variants that have been reported in disease-associated genes.
CRISPR-Cas9 editing	A method that uses an RNA guide and a DNA-binding protein to cleave DNA at a specific location to create sequence-specific changes via homologous recombination with a donor template.
Intestinal organoids	Epithelial "mini-guts" grown <i>in vitro</i> from biopsies of the rectal mucosa or from stem cells from a single individual.
Airway submucosal glands	Mucus-secreting glands found in the connective tissue that provide fluid for hydrating the surface of the airway epithelial cells and enabling ciliary function.
Airway surface	Liquid Fluid interface between the air and the cells in the lungs that confers protection from infection and facilitates removal of foreign particles.
Tracheal rings	Incomplete rings of highly elastic cartilage found in the anterior two-thirds of the tracheal wall.
Endocrine pancreas	Portion of the pancreas that produces hormones (insulin and glucagon) that are essential for glucose homeostasis.
Pseudomonas aeruginosa	Widely distributed gram-negative bacteria that show a predilection for acute and chronic infection of the lungs of individuals with cystic fibrosis.
Meconium ileus	Obstruction of the gut that usually develops <i>in utero</i> in the ileum of the small intestine and that is highly suggestive of cystic fibrosis.
Airway flow measurements	Series of standardized tests assessing the rate and volume of air that can be inhaled and exhaled: they are used to determine the degree of disease in the lungs in individuals with cystic fibrosis.

Vas deferens	A tubular structure that conveys sperm from the testis to the urethra of the penis.
Disseminated bronchiectasis	Persistent dilation of the airways (bronchi) throughout the lungs.
Phase III clinical trials	The third of four phases of evaluating a drug in affected subjects that confirms its safety and efficacy.
Nasal potential difference	Measurement of voltage across nasal epithelium that represents the transport of ions and that, under specific conditions, can assess the function of cystic fibrosis transmembrane conductance regulator (CFTR) <i>in vivo</i> .
Open probability	A measure of the average fraction of time that a channel is open.
Phase II clinical trial	The second of four phases of evaluating a drug in affected subjects that establishes the efficacy of a drug compared to a placebo.
Theratypes	A recently invented term used to classify disease-associated DNA variants according to the molecular-based treatment to which they respond.

References

- MacKenzie T, et al. Longevity of patients with cystic fibrosis in 2000 to 2010 and beyond: survival analysis of the cystic fibrosis foundation patient registry. *Ann. Intern. Med.* 2014; 161:233–241. [PubMed: 25133359]
- Cystic Fibrosis Foundation. Cystic Fibrosis Foundation Patient Registry Annual Data Report 2011. Cystic Fibrosis Foundation; 2012.
- Rommens JM, et al. Identification of the cystic fibrosis gene: chromosome walking and jumping. *Science.* 1989; 245:1059–1065. [PubMed: 2772657]
- Riordan JR, et al. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science.* 1989; 245:1066–1073. [PubMed: 2475911]
- Kerem B, et al. Identification of the cystic fibrosis gene: genetic analysis. *Science.* 1989; 245:1073–1080. References 3–5 are landmark papers from 25 years ago reporting the discovery of the CFTR gene. [PubMed: 2570460]
- Kartner N, et al. Expression of the cystic fibrosis gene in non-epithelial invertebrate cells produces a regulated anion conductance. *Cell.* 1991; 64:681–691. [PubMed: 1705179]
- Quinton PM. Chloride impermeability in cystic fibrosis. *Nature.* 1983; 301:421–422. [PubMed: 6823316]
- Knowles MR, et al. Abnormal ion permeation through cystic fibrosis respiratory epithelium. *Science.* 1983; 221:1067–1070. [PubMed: 6308769]
- Welsh MJ, Smith AE. Molecular mechanisms of CFTR chloride channel dysfunction in cystic fibrosis. *Cell.* 1993; 73:1251–1254. This seminal review proposes a classification of variants based on their predominant effect on CFTR processing or function. [PubMed: 7686820]
- Rich DP, et al. Expression of cystic fibrosis transmembrane conductance regulator corrects defective chloride channel regulation in cystic fibrosis airway epithelial cells. *Nature.* 1990; 347:358–363. [PubMed: 1699126]
- Guggino WB, Stanton BA. New insights into cystic fibrosis: molecular switches that regulate CFTR. *Nature Rev. Mol. Cell. Biol.* 2006; 7:426–436. [PubMed: 16723978]

12. Moskowitz SM, et al. Clinical practice and genetic counseling for cystic fibrosis and CFTR-related disorders. *Genet. Med.* 2008; 10:851–868. [PubMed: 19092437]
13. Zielenski J. Genotype and phenotype in cystic fibrosis. *Respiration.* 2000; 67:117–133. [PubMed: 10773783]
14. Bombieri C, et al. Recommendations for the classification of diseases as CFTR-related disorders. *J. Cyst. Fibros.* 2011; 10(Suppl. 2):S86–S102. [PubMed: 21658649]
15. Ramsey BW, et al. A CFTR potentiator in patients with cystic fibrosis and the G551D mutation. *N. Engl. J. Med.* 2011; 365:1663–1672. [PubMed: 22047557]
16. Boyle MP, et al. A CFTR corrector (lumacaftor) and a CFTR potentiator (ivacaftor) for treatment of patients with cystic fibrosis who have a phe508del CFTR mutation: a phase 2 randomised controlled trial. *Lancet Respir. Med.* 2014; 2:527–538. [PubMed: 24973281]
17. Cheng SH, et al. Defective intracellular transport and processing of CFTR is the molecular basis of most cystic fibrosis. *Cell.* 1990; 63:827–834. [PubMed: 1699669]
18. Denning GM, et al. Processing of mutant cystic fibrosis transmembrane conductance regulator is temperature-sensitive. *Nature.* 1992; 358:761–764. [PubMed: 1380673]
19. Lazrak A, et al. The silent codon change I507-ATC.ATT contributes to the severity of the .F508 CFTR channel dysfunction. *FASEB J.* 2013; 27:4630–4645. [PubMed: 23907436]
20. Sosnay PR, et al. Defining the disease liability of variants in the cystic fibrosis transmembrane conductance regulator gene. *Nature Genet.* 2013; 45:1160–1167. [PubMed: 23974870]
21. Van Goor F, Yu H, Burton B, Hoffman BJ. Effect of ivacaftor on CFTR forms with missense mutations associated with defects in protein processing or function. *J. Cyst. Fibros.* 2014; 13:29–36. [PubMed: 23891399]
22. Lukacs GL, Verkman AS. CFTR: folding, misfolding and correcting the F508 conformational defect. *Trends Mol. Med.* 2012; 18:81–91. [PubMed: 22138491]
23. Seibert FS, et al. Disease-associated mutations in the fourth cytoplasmic loop of cystic fibrosis transmembrane conductance regulator compromise biosynthetic processing and chloride channel activity. *J. Biol. Chem.* 1996; 271:15139–15145. [PubMed: 8662892]
24. Cotten JF, Ostedgaard LS, Carson MR, Welsh MJ. Effect of cystic fibrosis-associated mutations in the fourth intracellular loop of cystic fibrosis transmembrane conductance regulator. *J. Biol. Chem.* 1996; 271:21279–21284. [PubMed: 8702904]
25. Mendoza JL, Thomas PJ. Building an understanding of cystic fibrosis on the foundation of ABC transporter structures. *J. Bioenerg. Biomembr.* 2007; 39:499–505. [PubMed: 18080175]
26. Mornon JP, Lehn P, Callebaut I. Atomic model of human cystic fibrosis transmembrane conductance regulator: membrane-spanning domains and coupling interfaces. *Cell. Mol. Life Sci.* 2008; 65:2594–2612. [PubMed: 18597042]
27. Serohijos AW, et al. Phenylalanine-508 mediates a cytoplasmic-membrane domain contact in the CFTR 3D structure crucial to assembly and channel function. *Proc. Natl Acad. Sci. USA.* 2008; 105:3256–3261. References 25–27 provide a foundation for the three-dimensional modelling of CFTR. [PubMed: 18305154]
28. Thibodeau PH, et al. The cystic fibrosis-causing mutation F508 affects multiple steps in cystic fibrosis transmembrane conductance regulator biogenesis. *J. Biol. Chem.* 2010; 285:35825–35835. [PubMed: 20667826]
29. Krasnov KV, Tzetis M, Cheng J, Guggino WB, Cutting GR. Localization studies of rare missense mutations in cystic fibrosis transmembrane conductance regulator (CFTR) facilitate interpretation of genotype-phenotype relationships. *Hum. Mutat.* 2008; 29:1364–1372. [PubMed: 18951463]
30. Rabe WM, et al. Correction of both NBD1 energetics and domain interface is required to restore F508 CFTR folding and function. *Cell.* 2012; 148:150–163. [PubMed: 22265408]
31. Mendoza JL, et al. Requirements for efficient correction of F508 CFTR revealed by analyses of evolved sequences. *Cell.* 2012; 148:164–174. [PubMed: 22265409]
32. Okiyoneda T, et al. Mechanism-based corrector combination restores F508-CFTR folding and function. *Nature Chem. Biol.* 2013; 9:444–454. [PubMed: 23666117]
33. Matsui H, et al. Evidence for periciliary liquid layer depletion, not abnormal ion composition, in the pathogenesis of cystic fibrosis airways disease. *Cell.* 1998; 95:1005–1015. [PubMed: 9875854]

34. Wong AP, et al. Directed differentiation of human pluripotent stem cells into mature airway epithelia expressing functional CFTR protein. *Nature Biotech.* 2012; 30:876–882.
35. Dekkers JF, et al. A functional CFTR assay using primary cystic fibrosis intestinal organoids. *Nature Med.* 2013; 19:939–945. [PubMed: 23727931]
36. Schwank G, et al. Functional repair of CFTR by CRISPR/Cas9 in intestinal stem cell organoids of cystic fibrosis patients. *Cell Stem Cell.* 2013; 13:653–658. [PubMed: 24315439]
37. Supryniewicz FA, et al. Conditionally reprogrammed cells represent a stem-like state of adult epithelial cells. *Proc. Natl Acad. Sci. USA.* 2012; 109:20035–20040. [PubMed: 23169653]
38. Snouwaert JN, et al. An animal model for cystic fibrosis made by gene targeting. *Science.* 1992; 257:1083–1088. [PubMed: 1380723]
39. Tuggle KL, et al. Characterization of defects in ion transport and tissue development in cystic fibrosis transmembrane conductance regulator (CFTR)-knockout rats. *PLoS ONE.* 2014; 9:e91253. [PubMed: 24608905]
40. Sun X, et al. Disease phenotype of a ferret CFTR knockout model of cystic fibrosis. *J. Clin. Invest.* 2010; 120:3149–3160. [PubMed: 20739752]
41. Rogers CS, et al. Disruption of the CFTR gene produces a model of cystic fibrosis in newborn pigs. *Science.* 2008; 321:1837–1841. References 40 and 41 are the first reports to describe the phenotypes of ferrets and pigs caused by loss of CFTR function. [PubMed: 18818360]
42. Navis A, Marjoram L, Bagnat M. Cfr controls lumen expansion and function of Kupffer's vesicle in zebrafish. *Development.* 2013; 140:1703–1712. [PubMed: 23487313]
43. Wilke M, et al. Mouse models of cystic fibrosis: phenotypic analysis and research applications. *J. Cyst. Fibros.* 2011; 10:S152–S171. [PubMed: 21658634]
44. Grubb BR, Boucher RC. Pathophysiology of gene-targeted mouse models for cystic fibrosis. *Physiol. Rev.* 1999; 79:S193–S214. [PubMed: 9922382]
45. Clarke LB, et al. Relationship of a non-cystic fibrosis transmembrane conductance regulator-mediated chloride conductance to organ-level disease in CFTR^{-/-} mice. *Proc. Natl Acad. Sci. USA.* 1994; 91:479–483. [PubMed: 7507247]
46. Rock JR, et al. Transmembrane protein 16A (TMEM16A) is a Ca²⁺-regulated Cl⁻ secretory channel in mouse airways. *J. Biol. Chem.* 2009; 284:14875–14880.
47. Henderson LB, et al. Variation in MSRA modifies risk of neonatal intestinal obstruction in cystic fibrosis. *PLoS Genet.* 2012; 8:e1002580. [PubMed: 22438829]
48. Keiser NW, Engelhardt JF. New animal models of cystic fibrosis: what are they teaching us? *Curr. Opin. Pulm. Med.* 2011; 17:478–483. [PubMed: 21857224]
49. Rogers CS, et al. Production of CFTR-null and CFTR- F508 heterozygous pigs by adeno-associated virus-mediated gene targeting and somatic cell nuclear transfer. *J. Clin. Invest.* 2008; 118:1571–1577. [PubMed: 18324337]
50. Stoltz DA, et al. Cystic fibrosis pigs develop lung disease and exhibit defective bacterial eradication at birth. *Sci. Transl. Med.* 2010; 2:29ra31.
51. Khan TZ, et al. Early pulmonary inflammation in infants with cystic fibrosis. *Am. J. Respir. Crit. Care Med.* 1995; 151:1075–1082. [PubMed: 7697234]
52. Pezzulo AA, et al. Reduced airway surface pH impairs bacterial killing in the porcine cystic fibrosis lung. *Nature.* 2012; 487:109–113. [PubMed: 22763554]
53. Adam RJ, et al. Air trapping and airflow obstruction in newborn cystic fibrosis piglets. *Am. J. Respir. Crit. Care Med.* 2013; 188:1434–1441. [PubMed: 24168209]
54. Chen JH, et al. Loss of anion transport without increased sodium absorption characterizes newborn porcine cystic fibrosis airway epithelia. *Cell.* 2010; 143:911–923. [PubMed: 21145458]
55. Hodges CA, Grady BR, Mishra K, Cotton CU, Drumm ML. Cystic fibrosis growth retardation is not correlated with loss of Cfr in the intestinal epithelium. *Am. J. Physiol. Gastrointest. Liver Physiol.* 2011; 301:G528–G536. [PubMed: 21659619]
56. Stoltz DA, et al. Intestinal CFTR expression alleviates meconium ileus in cystic fibrosis pigs. *J. Clin. Invest.* 2013; 123:2685–2693. [PubMed: 23676501]
57. Ode KL, Moran A. New insights into cystic fibrosis-related diabetes in children. *Lancet Diabetes Endocrinol.* 2013; 1:52–58. [PubMed: 24622267]

58. Olivier AK, et al. Abnormal endocrine pancreas function at birth in cystic fibrosis ferrets. *J. Clin. Invest.* 2012; 122:3755–3768. [PubMed: 22996690]
59. Uc A, et al. Glycaemic regulation and insulin secretion are abnormal in cystic fibrosis pigs despite sparing of islet cell mass. *Clin. Sci. (Lond.)*. 2015; 128:131–142. [PubMed: 25142104]
60. Kerem E, et al. The relation between genotype and phenotype in cystic fibrosis--analysis of the most common mutation (.F508). *N. Engl. J. Med.* 1990; 323:1517–1522. [PubMed: 2233932]
61. The Cystic Fibrosis Genotype-Phenotype Consortium. Correlation between genotype and phenotype in patients with cystic fibrosis. *N. Engl. J. Med.* 1993; 329:1308–1313. [PubMed: 8166795]
62. Wilschanski M, et al. Correlation of sweat chloride concentration with classes of the cystic fibrosis transmembrane conductance regulator gene mutations. *J. Pediatr.* 1995; 127:705–710. [PubMed: 7472820]
63. McKone EF, Goss CH, Aitken ML. CFTR genotype as a predictor of prognosis in cystic fibrosis. *Chest.* 2006; 130:1441–1447. [PubMed: 17099022]
64. Gan K-H, et al. A cystic fibrosis mutation associated with mild lung disease. *N. Engl. J. Med.* 1995; 333:95–99. [PubMed: 7539891]
65. Mekus F, et al. Categories of .F508 homozygous cystic fibrosis twin and sibling pairs with distinct phenotypic characteristics. *Twin. Res.* 2000; 3:277–293. [PubMed: 11463149]
66. Vanscoy LL, et al. Heritability of lung disease severity in cystic fibrosis. *Am. J. Respir. Crit. Care Med.* 2007; 175:1036–1043. [PubMed: 17332481]
67. Collaco JM, Blackman SM, McGready J, Naughton KM, Cutting GR. Quantification of the relative contribution of environmental and genetic factors to variation in cystic fibrosis lung function. *J. Pediatr.* 2010; 157:802–807. [PubMed: 20580019]
68. Li Z, et al. Longitudinal development of mucoid *Pseudomonas aeruginosa* infection and lung disease progression in children with cystic fibrosis. *JAMA.* 2005; 293:581–588. [PubMed: 15687313]
69. Green DM, et al. Heritability of respiratory infection with *Pseudomonas aeruginosa* in cystic fibrosis. *J. Pediatr.* 2012; 161:290–295. [PubMed: 22364820]
70. Bradley GM, Blackman SM, Watson CP, Doshi VK, Cutting GR. Genetic modifiers of nutritional status in cystic fibrosis. *Am. J. Clin. Nutr.* 2012; 96:1299–1308. [PubMed: 23134884]
71. Stanke F, et al. Genes that determine immunology and inflammation modify the basic defect of impaired ion conductance in cystic fibrosis epithelia. *J. Med. Genet.* 2011; 48:24–31. [PubMed: 20837493]
72. Blackman SM, et al. Genetic modifiers play a substantial role in diabetes complicating cystic fibrosis. *J. Clin. Endocrinol. Metab.* 2009; 94:1302–1309. [PubMed: 19126627]
73. Finkelstein SM, et al. Diabetes mellitus associated with cystic fibrosis. *J. Pediatr.* 1988; 112:373–377. [PubMed: 3346774]
74. Rozmahel R, et al. Modulation of disease severity in cystic fibrosis transmembrane conductance regulator deficient mice by a secondary genetic factor. *Nature Genet.* 1996; 12:280–287. [PubMed: 8589719]
75. Blackman SM, et al. Relative contribution of genetic and nongenetic modifiers to intestinal obstruction in cystic fibrosis. *Gastroenterology.* 2006; 131:1030–1039. [PubMed: 17030173]
76. Weiler CA, Drumm ML. Genetic influences on cystic fibrosis lung disease severity. *Front. Pharmacol.* 2013; 4:40. [PubMed: 23630497]
77. Cutting GR. Modifier genes in Mendelian disorders: the example of cystic fibrosis. *Ann. NY Acad. Sci.* 2010; 1214:57–69. [PubMed: 21175684]
78. Knowles MR, Drumm M. The influence of genetics on cystic fibrosis phenotypes. *Cold Spring Harb. Perspect. Med.* 2012; 2:a009548. [PubMed: 23209180]
79. Drumm ML, Ziady AG, Davis PB. Genetic variation and clinical heterogeneity in cystic fibrosis. *Annu. Rev. Pathol.* 2012; 7:267–282. [PubMed: 22017581]
80. Guillot L, et al. Lung disease modifier genes in cystic fibrosis. *Int. J. Biochem. Cell Biol.* 2014; 52:83–93. [PubMed: 24569122]

81. Drumm ML, et al. Gene modifiers of lung disease in cystic fibrosis. *N. Engl. J. Med.* 2005; 353:1443–1453. This is an outstanding example of a candidate gene association study in which TGFB1 was identified as a modifier of lung disease in cystic fibrosis. [PubMed: 16207846]
82. Bremer LA, et al. Interaction between a novel TGFB1 haplotype and CFTR genotype is associated with improved lung function in cystic fibrosis. *Hum. Mol. Genet.* 2008; 17:2228–2237. [PubMed: 18424453]
83. Chalmers JD, Fleming GB, Hill AT, Kilpatrick DC. Impact of mannose-binding lectin insufficiency on the course of cystic fibrosis: a review and meta-analysis. *Glycobiology.* 2011; 21:271–282. [PubMed: 21045008]
84. Dorfman R, et al. Complex two-gene modulation of lung disease severity in children with cystic fibrosis. *J. Clin. Invest.* 2008; 118:1040–1049. [PubMed: 18292811]
85. Collaco JM, et al. Interactions between secondhand smoke and genes that affect cystic fibrosis lung disease. *JAMA.* 2008; 299:417–424. [PubMed: 18230779]
86. Wright FA, et al. Genome-wide association and linkage identify modifier loci of lung disease severity in cystic fibrosis at 11p13 and 20q13.2. *Nature Genet.* 2011; 43:539–546. This paper demonstrates the successful application of genome-wide methods to the search for gene modifiers of cystic fibrosis. [PubMed: 21602797]
87. Blackman SM, et al. Genetic modifiers of cystic fibrosis-related diabetes. *Diabetes.* 2013; 62:3627–3635. [PubMed: 23670970]
88. Bartlett JR, et al. Genetic modifiers of liver disease in cystic fibrosis. *JAMA.* 2009; 302:1076–1083. [PubMed: 19738092]
89. Sun L, et al. Multiple apical plasma membrane constituents are associated with susceptibility to meconium ileus in individuals with cystic fibrosis. *Nature Genet.* 2012; 44:562–569. [PubMed: 22466613]
90. Li W, et al. Unraveling the complex genetic model for cystic fibrosis: pleiotropic effects of modifier genes on early cystic fibrosis-related morbidities. *Hum. Genet.* 2014; 133:151–161. [PubMed: 24057835]
91. Emond MJ, et al. Exome sequencing of extreme phenotypes identifies DCTN4 as a modifier of chronic *Pseudomonas aeruginosa* infection in cystic fibrosis. *Nature Genet.* 2012; 44:886–889. [PubMed: 22772370]
92. Luciani A, et al. Defective CFTR induces aggresome formation and lung inflammation in cystic fibrosis through ROS-mediated autophagy inhibition. *Nature Cell Biol.* 2010; 12:863–875. [PubMed: 20711182]
93. Grody WW, et al. Laboratory standards and guidelines for population-based cystic fibrosis carrier screening. *Genet. Med.* 2001; 3:149–154. [PubMed: 11280952]
94. Grody WW, Cutting GR, Watson MS. The cystic fibrosis mutation “arms race”: when less is more. *Genet. Med.* 2007; 9:739–744. [PubMed: 18007142]
95. Kelley TJ, Al Nakkash L, Cotton CU, Drumm ML. Activation of endogenous .F508 cystic fibrosis transmembrane conductance regulator by phosphodiesterase inhibition. *J. Clin. Invest.* 1996; 98:513–520. [PubMed: 8755664]
96. Van Goor F, et al. Rescue of CF airway epithelial cell function in vitro by a CFTR potentiator, VX-770. *Proc. Natl Acad. Sci. USA.* 2009; 106:18825–18830. This is the first report that ivacaftor potentiates the function of CFTR bearing the G551D mutation. [PubMed: 19846789]
97. Jih KY, Hwang TC. Vx-770 potentiates CFTR function by promoting decoupling between the gating cycle and ATP hydrolysis cycle. *Proc. Natl Acad. Sci. USA.* 2013; 110:4404–4409. [PubMed: 23440202]
98. Eckford PD, Li C, Ramjeesingh M, Bear CE. Cystic fibrosis transmembrane conductance regulator (CFTR) potentiator VX-770 (ivacaftor) opens the defective channel gate of mutant CFTR in a phosphorylation-dependent but ATP-independent manner. *J. Biol. Chem.* 2012; 287:36639–36649. [PubMed: 22942289]
99. Accurso FJ, et al. Effect of VX-770 in persons with cystic fibrosis and the G551D-CFTR mutation. *N. Engl. J. Med.* 2010; 363:1991–2003. References 15 and 99 are the first reports of a clinically effective treatment for cystic fibrosis based on targeting the molecular defect. [PubMed: 21083385]

100. Char JE, et al. A little CFTR goes a long way: CFTR-dependent sweat secretion from G551D and R117H-5T cystic fibrosis subjects taking ivacaftor. *PLoS ONE*. 2014; 9:e88564. [PubMed: 24520399]
101. Accurso FJ, et al. Sweat chloride as a biomarker of CFTR activity: proof of concept and ivacaftor clinical trial data. *J. Cyst. Fibros.* 2014; 13:139–147. [PubMed: 24660233]
102. Van Goor F, et al. Correction of the F508del-CFTR protein processing defect in vitro by the investigational drug VX-809. *Proc. Natl Acad. Sci. USA*. 2011; 108:18843–18848. [PubMed: 21976485]
103. Clancy JP, et al. Results of a phase IIa study of VX-809, an investigational CFTR corrector compound, in subjects with cystic fibrosis homozygous for the F508del-CFTR mutation. *Thorax*. 2012; 67:12–18. [PubMed: 21825083]
104. Cholon DM, et al. Potentiator ivacaftor abrogates pharmacological correction of .F508 CFTR in cystic fibrosis. *Sci. Transl Med.* 2014; 6:246ra96.
105. Veit G, et al. Some gating potentiators, including VX-770, diminish .F508-CFTR functional expression. *Sci. Transl Med.* 2014; 6:246ra97.
106. Phuan PW, et al. Synergy-based small-molecule screen using a human lung epithelial cell line yields . 508-CFTR correctors that augment VX-809 maximal efficacy. *Mol. Pharmacol.* 2014; 86:42–51.
107. De Boeck K, et al. CFTR biomarkers: time for promotion to surrogate end-point. *Eur. Respir. J.* 2013; 41:203–216. [PubMed: 22878883]
108. Yu H, et al. Ivacaftor potentiation of multiple CFTR channels with gating mutations. *J. Cyst. Fibros.* 2012; 11:237–245. [PubMed: 22293084]
109. Dietz HC. New therapeutic approaches to mendelian disorders. *N. Engl. J. Med.* 2010; 363:852–863. [PubMed: 20818846]
110. Mendell JT, Sharifi NA, Meyers JL, Martinez-Murillo F, Dietz HC. Nonsense surveillance regulates expression of diverse classes of mammalian transcripts and mutes genomic noise. *Nature Genet.* 2004; 36:1073–1078. [PubMed: 15448691]
111. Howard M, Frizzell RA, Bedwell DM. Aminoglycoside antibiotics restore CFTR function by overcoming premature stop mutations. *Nature Med.* 1996; 2:467–469. [PubMed: 8597960]
112. Kerem E, et al. Effectiveness of PTC124 treatment of cystic fibrosis caused by nonsense mutations: a prospective phase II trial. *Lancet.* 2008; 372:719–727. [PubMed: 18722008]
113. Kerem E, et al. Ataluren for the treatment of nonsense-mutation cystic fibrosis: a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet Respir. Med.* 2014; 2:539–547. [PubMed: 24836205]
114. Oren YS, et al. The unfolded protein response affects readthrough of premature termination codons. *EMBO Mol. Med.* 2014; 6:685–701. [PubMed: 24705877]
115. Havens MA, Duelli DM, Hastings ML. Targeting RNA splicing for disease therapy. *Wiley. Interdiscip. Rev. RNA.* 2013; 4:247–266. [PubMed: 23512601]
116. Griesenbach U, Alton EW. Moving forward: cystic fibrosis gene therapy. *Hum. Mol. Genet.* 2013; 22:R52–R58. [PubMed: 23918661]
117. Falconer, DS.; Mackay, TF. C. Heritability in *Introduction to Quantitative Genetics*. Pearson Education Limited; 1996. p. 160-183.
118. Anguiano A, et al. Congenital bilateral absence of the vas deferens - a primarily genital form of cystic fibrosis. *JAMA.* 1992; 267:1794–1797. [PubMed: 1545465]
119. Gilljam M, et al. Airway inflammation and infection in congenital bilateral absence of the vas deferens. *Am. J. Respir. Crit. Care Med.* 2003; 169:174–179. [PubMed: 14551163]
120. Colin AA, et al. Pulmonary function and clinical observations in men with congenital bilateral absence of the vas deferens. *Chest.* 1996; 110:440–445. [PubMed: 8697849]
121. Chillón M, et al. Mutations in the cystic fibrosis gene in patients with congenital absence of the vas deferens. *N. Engl. J. Med.* 1995; 332:1475–1480. [PubMed: 7739684]
122. Noone PG, Knowles MR. ‘CFTR-opathies’: disease phenotypes associated with cystic fibrosis transmembrane regulator gene mutations. *Respir. Res.* 2001; 2:328–332. [PubMed: 11737931]

123. Ooi CY, et al. Type of CFTR mutation determines risk of pancreatitis in patients with cystic fibrosis. *Gastroenterology*. 2011; 140:153–161. [PubMed: 20923678]
124. Larusch J, Solomon S, Whitcomb DC. Pancreatitis Overview. 2014 [online], <http://www.ncbi.nlm.nih.gov/books/NBK190101/>.
125. Larusch J, et al. Mechanisms of CFTR functional variants that impair regulated bicarbonate permeation and increase risk for pancreatitis but not for cystic fibrosis. *PLoS Genet*. 2014; 10:e1004376. [PubMed: 25033378]
126. Park HW, et al. Dynamic regulation of CFTR bicarbonate permeability by [Cl⁻]_i and its role in pancreatic bicarbonate secretion. *Gastroenterology*. 2010; 139:620–631. [PubMed: 20398666]
127. Kim RD, et al. Pulmonary nontuberculous mycobacterial disease: prospective study of a distinct preexisting syndrome. *Am. J. Respir. Crit. Care Med*. 2008; 178:1066–1074. [PubMed: 18703788]
128. Knutsen AP, Slavin RG. Allergic bronchopulmonary aspergillosis in asthma and cystic fibrosis. *Clin. Dev. Immunol*. 2011; 2011:843763. [PubMed: 21603163]
129. Wang X, et al. Mutation in the gene responsible for cystic fibrosis and predisposition to chronic rhinosinusitis in the general population. *JAMA*. 2000; 284:1814–1819. [PubMed: 11025834]
130. Wang XJ, Kim J, McWilliams R, Cutting GR. Increased prevalence of chronic rhinosinusitis in carriers of a cystic fibrosis mutation. *Arch. Otolaryngol. Head Neck Surg*. 2005; 131:237–240. [PubMed: 15781764]
131. Ooi CY, et al. Does extensive genotyping and nasal potential difference testing clarify the diagnosis of cystic fibrosis among patients with single-organ manifestations of cystic fibrosis? *Thorax*. 2014; 69:254–260. [PubMed: 24149827]
132. Cutting GR. Annotating DNA variants is the next major goal for human genetics. *Am. J. Hum. Genet*. 2014; 94:5–10. [PubMed: 24387988]
133. Amaral MD. Processing of CFTR: traversing the cellular maze - how much CFTR needs to go through to avoid cystic fibrosis? *Pediatr. Pulmonol*. 2005; 39:479–491. [PubMed: 15765539]
134. Highsmith WE Jr, et al. Identification of a splice site mutation (2789 + 5G>A) associated with small amounts of normal CFTR mRNA and mild cystic fibrosis. *Hum. Mutat*. 1997; 9:332–338. [PubMed: 9101293]
135. Ramalho AS, et al. Five percent of normal cystic fibrosis transmembrane conductance regulator mRNA ameliorates the severity of pulmonary disease in cystic fibrosis. *Am. J. Respir. Cell. Mol. Biol*. 2002; 27:619–627. [PubMed: 12397022]
136. Chu C-S, Trapnell BC, Curristin SM, Cutting GR, Crystal RG. Extensive posttranslational deletion of the coding sequences for part of nucleotide-binding fold 1 in respiratory epithelial mRNA transcripts of the cystic fibrosis transmembrane conductance regulator gene is not associated with the clinical manifestations of cystic fibrosis. *J. Clin. Invest*. 1992; 90:785–790. [PubMed: 1381723]
137. Rave-Harel N, et al. The molecular basis of partial penetrance of splicing mutations in cystic fibrosis. *Am. J. Hum. Genet*. 1997; 60:87–94. [PubMed: 8981951]
138. Johnson LG, et al. Efficiency of gene transfer for restoration of normal airway epithelial function in cystic fibrosis. *Nature Genet*. 1993; 2:21–25. [PubMed: 1284642]
139. Zhang L, et al. CFTR delivery to 25% of surface epithelial cells restores normal rates of mucus transport to human cystic fibrosis airway epithelium. *PLoS. Biol*. 2009; 7:e1000155. [PubMed: 19621064]
140. Romey MC, et al. A naturally occurring sequence variation that creates a YY1 element is associated with increased cystic fibrosis transmembrane conductance regulator gene expression. *J. Biol. Chem*. 2000; 275:3561–3567. [PubMed: 10652351]
141. Highsmith WE Jr, et al. A novel mutation in the cystic fibrosis gene in patients with pulmonary disease but normal sweat chloride concentrations. *N. Engl. J. Med*. 1994; 331:974–980. [PubMed: 7521937]
142. Hamosh A, Rosenstein BJ, Cutting GR. CFTR nonsense mutations G542X and W1282X associated with severe reduction of CFTR mRNA in nasal epithelial cells. *Hum. Mol. Genet*. 1992; 1:542–544. [PubMed: 1284888]

143. Silvis MR, et al. A mutation in the cystic fibrosis transmembrane conductance regulator generates a novel internalization sequence and enhances endocytic rates. *J. Biol. Chem.* 2003; 278:11554–11560. [PubMed: 12529365]
144. Wang Y, Wrennall JA, Cai Z, Li H, Sheppard DN. Understanding how cystic fibrosis mutations disrupt CFTR function: from single molecules to animal models. *Int. J. Biochem. Cell Biol.* 2014; 52:47–57. [PubMed: 24727426]
145. Friedman KJ, et al. Correction of aberrant splicing of the cystic fibrosis transmembrane conductance regulator (CFTR) gene by antisense oligonucleotides. *J. Biol. Chem.* 1999; 274:36193–36199. [PubMed: 10593905]
146. Xue X, et al. Synthetic aminoglycosides efficiently suppress cystic fibrosis transmembrane conductance regulator nonsense mutations and are enhanced by ivacaftor. *Am. J. Respir. Cell Mol. Biol.* 2014; 50:805–816. [PubMed: 24251786]
147. Meyerholz DK, et al. Loss of cystic fibrosis transmembrane conductance regulator function produces abnormalities in tracheal development in neonatal pigs and young children. *Am. J. Respir. Crit. Care Med.* 2010; 182:1251–1261. [PubMed: 20622026]




Figure 1. Cardinal features of cystic fibrosis and relative contribution of genetic modifiers to variation in select cystic fibrosis traits

A diagnosis of cystic fibrosis is based on the presence of clinical findings shown on the left, along with an elevated sweat chloride concentration (>60mM). The degree of organ system dysfunction varies considerably among affected individuals. Genetic modifiers and non-genetic factors both contribute to airway obstruction and infection with *Pseudomonas aeruginosa* -two traits that define lung disease in cystic fibrosis. Cystic fibrosis transmembrane conductance regulator (*CFTR*) genotype is the primary determinant of the degree of pancreatic exocrine dysfunction. The presence of *CFTR* variants associated with severe pancreatic exocrine dysfunction is essentially a pre-requisite for the development of diabetes and intestinal obstruction. In the setting of severe endocrine dysfunction, genetic modifiers determine when, and if, diabetes occurs and whether neonatal intestinal obstruction occurs. Genetic variation plays the predominant part in nutritional status as assessed by body mass index (BMI)⁷⁰.




Figure 2. Molecular consequences of variants in CFTR

The degree to which epithelial ion transport is altered in an individual with cystic fibrosis is determined by the effect of each disease-causing variant on the quantity and the function of cystic fibrosis transmembrane conductance regulator (CFTR). The key steps of CFTR biogenesis in an epithelial cell are depicted. The membrane-spanning domains of CFTR are shown as red boxes, the two nucleotide-binding domains as yellow circles, and the regulatory domain as a blue circle. The quantity of CFTR protein in the apical cell membrane is a product of the amount of RNA transcribed, the efficiency of RNA splicing, the fraction of protein correctly folded and the stability of the protein in the membrane. The level and/or content of *CFTR* transcripts can be affected by disease-causing variants in the promoter (for example, c.-234T→A (also known as -102T→A in legacy nomenclature))¹⁴⁰ and splice sites (for example, c. 3717+12191 C→T (legacy 3849+10 kb C→1))¹⁴¹, or by variants that introduce a premature termination codon (PTC) and that lead to RNA decay (for example, p.Gly542X; (legacy G542X))¹⁴². The processing of CFTR can be altered by variants that cause aberrant folding of the protein, leading to degradation (for example, p.PheS08del (legacy F508del))¹⁸, or by variants that cause reduced membrane stability as a result of increased rates of endocytosis (for example, p.Asn287Tyr (legacy N287Y))¹⁴³. The function of CFTR is dependent on activity of the ion channel and on the efficiency of conductance of ions through the channel. Disease-causing variants cause reduction in activity (for example, p.Gly551Asp (legacy G551D))¹⁴⁴ or changes in the conductance properties of the chloride channel (for example, p.Arg334Trp (legacy R334W))¹⁴⁴. cAMP, cyclic AMP; ER, endoplasmic reticulum.




Figure 3. Molecular treatments for cystic fibrosis

The effects of various therapies (shown as arrows) on the quantity and/or function of mutant cystic fibrosis transmembrane conductance regulator (C FTR; FIG. 2) are shown. At least 10% of 'normal' CFTR function is required for 'clinical response' in lung function (grey region). The indistinct border of the clinical response accommodates various estimates of the amount of C FTR ion transport that have to be achieved and the probability that the product of quantity and function may generate a nonlinear function in certain circumstances. Red circles indicate the approximate product of quantity and function of CFTR bearing the indicated variant. Yellow circles indicate a shift in the product of quantity and function that does not achieve a clinical response, and green circles indicate a shift that does produce a clinical response.

a | Single-drug strategies for G551D (also known as p. Gly551Asp) and F508del (also known as p.Phe508del) are shown. CFTR bearing the G551D variant is found at normal levels in the cell membrane but cannot be activated. The potentiator ivacaftor increases the activity of G551D-C FTR, thereby increasing chloride transport in airway epithelia to a level that produces a clinical response in the lungs. The common variant F508del causes a defect in protein folding, which greatly diminishes CFTR quantity at the plasma membrane and reduces both membrane residency and channel activity. Application of lumacaftor to cells increases the quantity and, to a lesser degree, the chloride transport of F508del-CFTR, but not to a level that produces a clinical response in the lungs.

b | Combinatorial strategies for F508del are shown. In cell-based studies, chronic administration of ivacaftor counteracts the increase in CFTR stability conferred by lumacaftor^{119,120}. Use of other potentiators that do not antagonize the effect of lumacaftor, such as the investigational compound P5, could provide higher therapeutic benefit¹²⁰. An alternative approach is to combine correctors that affect different stages of CFTR folding, such as lumacaftor and C4 (REF. 32).

c | Therapy strategy for rare variants is shown. Red circles with asterisks indicate variants that permit production of normal or near-normal quantity of CFTR but that alter the activity of the C FTR chloride channel. As this functional defect is similar to that caused by G551D, each variant is tested for response to ivacaftor in cell-based studies. Recovery of C FTR function that exceeds 10% of levels seen in normal individuals indicates that a clinical response is expected, thereby justifying clinical trials.

d | Strategies for nonsense and splice site variants are shown. The splice site variant 3849+10kbC→T (also known as c.3717+12191C→T) activates a cryptic splice donor site, causing reduction of the full-length CFTR transcript to ~8% of normal levels. Suppression of the cryptic splice site can increase CFTR protein levels above 10% of normal¹⁴⁵. The nonsense variant G542X (also known as p.Gly542X) introduces a premature termination codon (PTC) that causes severe reduction in mRNA levels and an absence of the CFTR protein. Use of

PTC suppressors increases transcript and protein levels, leading to a modest recovery of CFTR function that falls short of a clinical response. Combining ivacaftor with a PTC suppressor produces a 2.5-4-fold increase in function of G542X-CFTR that could be sufficient to produce a clinical response in lung function¹⁴⁶.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1Phenotype features in humans and animal models with mutated *CFTR*

Features*	Human	Pig	Ferret	Mouse	Rat	Zebrafish
Aberrant chloride transport	↑	↑	↑	↑	↑	↑
Intestinal obstruction	↑	↑↑	↑↑	↑↑ [‡]	↑↑ [‡]	-
Growth disturbance	↑	↑↑	↑↑	↑↑	↑	-
Maldevelopment of trachea	↑	↑	↑	↑	↑	-
Pancreatic exocrine dysfunction	↑	↑↑	↑	-	-	-
Obstructive lung disease	↑	↑	↑	-	-	-
Liver dysfunction	↑	↑↑	↑	-	-	-
Diabetes mellitus	↑	↑	↑	-	-	-
Anomalous vas deferens	↑	↑	↑	-	↑	-

CFTR, cystic fibrosis transmembrane conductance regulator.* Data are summarized from reviews^{43,48} and primary publications^{39,42,58,59,147}.[‡] Intestinal obstruction occurs after the neonatal period in murine models around the time of weaning.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript