


Illuminating G-Protein-Coupling Selectivity of GPCRs

Graphical Abstract



Authors

Asuka Inoue, Francesco Raimondi, Francois Marie Ngako Kadji, ..., J. Silvio Gutkind, Junken Aoki, Robert B. Russell

Correspondence

iaska@tohoku.ac.jp (A.I.), francesco.raimondi@bioquant.uni-heidelberg.de (F.R.), robert.russell@bioquant.uni-heidelberg.de (R.B.R.)

In Brief

A large-scale functional interaction study reveals how GPCRs engage with subsets of G(alpha) proteins and outlines characteristic predictive of these associations.

Highlights

- Large datasets of quantitative coupling between 148 human GPCRs and 11 G proteins
- Identification of GPCR sequence-encoded features underlying G-protein selectivity
- A powerful predictor for scoring G-protein coupling from a given GPCR sequence
- Data-driven engineering of G12-selective designer GPCRs



Illuminating G-Protein-Coupling Selectivity of GPCRs

Asuka Inoue,^{1,2,3,7,8,*} Francesco Raimondi,^{4,5,7,*} Francois Marie Ngako Kadji,¹ Gurdeep Singh,^{4,5} Takayuki Kishi,¹ Akiharu Uwamizu,¹ Yuki Ono,¹ Yuji Shinjo,¹ Satoru Ishida,¹ Nadia Arang,⁶ Kouki Kawakami,¹ J. Silvio Gutkind,⁶ Junken Aoki,^{1,3} and Robert B. Russell^{4,5,*}

¹Graduate School of Pharmaceutical Sciences, Tohoku University, Sendai, Miyagi 980-8578, Japan

²Advanced Research & Development Programs for Medical Innovation (PRIME), Japan Agency for Medical Research and Development (AMED), Chiyoda-ku, Tokyo 100-0004, Japan

³Advanced Research & Development Programs for Medical Innovation (LEAP), AMED, Chiyoda-ku, Tokyo 100-0004, Japan

⁴CellNetworks, Bioquant, Heidelberg University, Im Neuenheimer Feld 267, 69120 Heidelberg, Germany

⁵Biochemie Zentrum Heidelberg (BZH), Heidelberg University, Im Neuenheimer Feld 328, 69120 Heidelberg, Germany

⁶Department of Pharmacology and Moores Cancer Center, University of California, San Diego, La Jolla, CA 92093, USA

⁷These authors contributed equally

⁸Lead Contact

*Correspondence: iaska@tohoku.ac.jp (A.I.), francesco.raimondi@bioquant.uni-heidelberg.de (F.R.), robert.russell@bioquant.uni-heidelberg.de (R.B.R.)

<https://doi.org/10.1016/j.cell.2019.04.044>

SUMMARY

Heterotrimeric G proteins consist of four subfamilies (G_s , $G_{i/o}$, $G_{q/11}$, and $G_{12/13}$) that mediate signaling via G-protein-coupled receptors (GPCRs), principally by receptors binding $G\alpha$ C termini. G-protein-coupling profiles govern GPCR-induced cellular responses, yet receptor sequence selectivity determinants remain elusive. Here, we systematically quantified ligand-induced interactions between 148 GPCRs and all 11 unique $G\alpha$ subunit C termini. For each receptor, we probed chimeric $G\alpha$ subunit activation via a transforming growth factor- α (TGF- α) shedding response in HEK293 cells lacking endogenous $G_{q/11}$ and $G_{12/13}$ proteins, and complemented G-protein-coupling profiles through a NanoBiT-G-protein dissociation assay. Interrogation of the dataset identified sequence-based coupling specificity features, inside and outside the transmembrane domain, which we used to develop a coupling predictor that outperforms previous methods. We used the predictor to engineer designer GPCRs selectively coupled to G_{12} . This dataset of fine-tuned signaling mechanisms for diverse GPCRs is a valuable resource for research in GPCR signaling.

INTRODUCTION

G-protein-coupled receptors (GPCRs), one of the largest protein superfamilies, are key mediators linking extracellular ligands to downstream signals and are the most common targets for pharmaceutical drug development (Hauser et al., 2017, 2018). Ligand binding induces conformational changes in GPCRs that then lead to intracellular binding by particular heterotrimeric G-protein

complexes, each consisting of $G\alpha$, $G\beta$, and $G\gamma$ subunits, where distinct $G\alpha$ subunits specify both GPCR interactions and the transduction of particular downstream signaling events (Wett- schureck and Offermanns, 2005). The human genome encodes 16 $G\alpha$ genes that are grouped into four subfamilies $G\alpha_s$, $G\alpha_{i/o}$, $G\alpha_{q/11}$, and $G\alpha_{12/13}$ that capture broad properties of downstream signaling (e.g., adenylyl cyclase activation by $G\alpha_s$) (Wett- schureck and Offermanns, 2005). In general, each of the hundreds of mammalian GPCRs couple with more than one G protein giving each a distinct coupling profile (Harding et al., 2018), or signature, which evokes a unique cellular response. Determining these GPCR profiles is critical to understanding their biology and pharmacology.

Pharmaceutical interest in GPCRs has prompted many efforts during the last decades to determine both their ligands and signaling (Hauser et al., 2018). Among approximately 360 non-sensory GPCR genes encoded in the human genome, one-third are still labeled as *orphans* to reflect the fact that either ligands and/or signaling are unknown (Harding et al., 2018). Previous efforts to uncover signaling profiles have been laborious and not standardized yet tended to identify only the subfamily-level signaling outcome (e.g., Ca^{2+} , cAMP, inositol phosphate, Rho activation), rather than the specific $G\alpha$ subunit binding event (Thomsen et al., 2005). Although this has led a collection of data on GPCR ligands and signaling exemplified in the IUPHAR/BPS Guide to Pharmacology (GtoPdb) (Harding et al., 2018), these databases have issues with mixed quality of G-protein-coupling data as well as the lack of “negative” coupling information. Certain G proteins are still comparatively understudied in terms of their GPCR partners, particularly $G_{12/13}$, which signal principally through Rho GTPases. Moreover, for the majority of well-studied receptors, only the *primary* (i.e., the most prominent) coupling is known, with *secondary* couplings known only for a minority. Yet, this G-protein-coupling information is limited to binary (primary coupling and not stated) or tertiary (primary, secondary couplings, and not stated) scoring and fails to provide






Figure 1. Chimeric G-Protein-Based TGF- α Shedding Assay to Probe Interaction between an Active GPCR and a C-Terminal Tail of a G α Subunit

(A) Mechanism of the TGF- α shedding assay. $G_{\alpha_{q/11}}$ - and/or $G_{12/13}$ -coupled receptors induce activation of a membrane-bound metalloprotease ADAM17, which is endogenously expressed in HEK293 cells, and subsequent ectodomain shedding of the alkaline phosphatase-fused TGF- α (AP-TGF- α) reporter construct. AP-TGF- α release into conditioned media is quantified through a colorimetric reaction. Parental HEK293 cells and cells devoid of the $G_{\alpha_{q/11}}$ subunits (ΔG_q), the $G_{\alpha_{12/13}}$ subunits (ΔG_{12}), or the $G_{\alpha_{q/11/12/13}}$ subunits ($\Delta G_q/\Delta G_{12}$) were used in the TGF- α shedding assay.

(B) Blunted TGF- α shedding response in the HEK293 cells devoid of the $G_{\alpha_{q/11}}$ and the $G_{12/13}$ subfamilies. GPCRs known to couple with $G_{12/13}$ (LPAR6 and PTGER3), $G_{\alpha_{q/11}}$ (CHRM1 and HRH1), and both (GALR2 and GHSR) were examined for ligand-induced TGF- α shedding responses in the parental HEK293 cells or the indicated G-protein-deficient cells. Symbols and error bars represent mean and SEM, respectively, of 3–6 independent experiments with each performed in triplicate.

(C) Chimeric G-protein-based TGF- α shedding assay in $\Delta G_q/\Delta G_{12}$ cells. A test GPCR is expressed in $\Delta G_q/\Delta G_{12}$ cells together with one of 11 chimeric G_{α} subunits harboring C-terminal 6-amino acid substitution and restoration of ligand-induced AP-TGF- α release response is measured. Note that there are 11 unique C-terminal sequences for the 16 human G_{α} subunits (the C-terminal 6-amino acid sequences of $G_{\alpha_{11}}$, $G_{\alpha_{12}}$, $G_{\alpha_{13}}$, $G_{\alpha_{14}}$, and $G_{\alpha_{15}}$ and those of G_{α_q} and $G_{\alpha_{11}}$ are identical; also see Figures S2A–S2C) and that the invariant leucine is encoded at the –7 position. The C-terminally truncated G_{α_q} construct ($G_{\alpha_q}(\Delta C)$) is used for a negative control.

(D) Representative data for the chimeric G-protein-based assay. TBXA2R was expressed with one of the 11 G_{α_q} constructs or the $G_{\alpha_q}(\Delta C)$ and treated with titrated concentration of a ligand (U-46619). AP-TGF- α release responses were fitted to a sigmoidal concentration-response curve (upper panels). G-protein

(legend continued on next page)

quantitative data sufficient to aid successful analyses, for example of GPCR residues involved in G-protein-coupling selectivity.

Efforts to predict coupling on the basis of sequence features have been made to complement the absence of a complete picture of G-protein signaling (Sgourakis et al., 2005b; Yabuki et al., 2005). For $G_{12/13}$, owing to limited availability of signaling assays, coupling information is incomplete. In addition, for orphan GPCRs, which lack pharmacological compounds to activate receptors, an accurate signaling prediction is desired to investigate not only coupling information, but also putative ligands. Although many methods have been employed, previous researches generally sought to identify broad sequence properties at particular sites on the sequences that are indicative of a particular coupling subgroup. These methods have met with mixed success and with exceptionally poor performances for $G_{12/13}$.

Recent structures of GPCRs in complex with G proteins (Carpenter et al., 2016; Draper-Joyce et al., 2018; García-Nafria et al., 2018a, 2018b; Kang et al., 2018; Koehl et al., 2018; Rasmussen et al., 2011) have revealed novel insights into G-protein coupling, including sequence determinants, and the flexibility of critical contact points (e.g., transmembrane helix 6, TM6) regulating G-protein access (Kang et al., 2018; Koehl et al., 2018). In particular, the C-terminal tail (Helix 5) of the $G\alpha$ subunit predominates at the interface with a GPCR, typically accounting for ~70% of the interacting surface (Flock et al., 2017; Rasmussen et al., 2011). This is consistent with earlier mutagenesis studies, which also highlight the importance of this region. Chimeric $G\alpha$ subunits, where 4–6 C-terminal residues are interchanged, can show swapped coupling selectivities: for example, the chimera $G\alpha_{q/11}$, with a $G\alpha_q$ backbone and a $G\alpha_{11}$ C terminus, has been used to monitor G_i signaling (Conklin et al., 1993).

Despite many advances in the understanding of GPCRs, the mechanisms by which they specifically signal through G proteins remain poorly understood (Koehl et al., 2018). This could be greatly enhanced by systematic screens of multiple, diverse GPCRs against a complete set of G proteins. Here, we exploited an experimentally derived dataset of GPCR/G-protein pairings, where we assessed coupling via chimeric G-protein-mediated responses of transforming growth factor- α (TGF- α) release. The extensive dataset provides new insights into the spectrum of coupling for 148 diverse receptors and enabled us to devise a highly accurate computational approach to predict coupling from sequence, which outperforms previous methods. The dataset, prediction method, and associated analysis provides new insights into features determining specificity, which we used to design the first G_{12} -specific engineered (chemogenetic) receptors that respond to a synthetic ligand.

RESULTS

$G_{q/11}$ - and $G_{12/13}$ -Dependent TGF- α Shedding Responses

To evaluate G-protein coupling, we exploited a TGF- α shedding assay (Figure 1A), which we showed previously to be a robust, high-throughput means to measure accumulated GPCR signals (Inoue et al., 2012). In the assay, we detect ADAM17-induced ectodomain shedding of alkaline phosphatase-fused TGF- α (AP-TGF- α) and subsequent release into conditioned media. We previously observed that $G_{q/11}$ - or $G_{12/13}$ -coupled receptors induce this process (Inoue et al., 2012), which we first tested using a panel of HEK293 cells lacking one or both of the $G_{q/11}$ and the $G_{12/13}$ subfamilies (hereafter denoted as ΔG_q , ΔG_{12} and $\Delta G_q/\Delta G_{12}$; Figure 1A) (Devost et al., 2017; Schrage et al., 2015). We tested GPCRs (Figure 1B) that are reported to couple with either $G_{q/11}$ (CHRM1 and HRH1 [Harding et al., 2018]) or $G_{12/13}$ (LPAR6 and PTGER3 [Kihara et al., 2014; Sugimoto and Narumiya, 2007]) or both (GALR2 and GHSR [Harding et al., 2018]). TGF- α shedding responses of the $G_{q/11}$ -coupled receptors and the $G_{12/13}$ -coupled receptors were diminished in ΔG_q and ΔG_{12} cells, respectively, while the responses were retained in cells lacking uncoupled G proteins. In the receptors coupling to both, the TGF- α shedding responses remained in ΔG_q and ΔG_{12} cells. For all tested GPCRs, TGF- α shedding responses were completely abolished in $\Delta G_q/\Delta G_{12}$ cells nor could G_s - or $G_{i/o}$ -coupled receptors induce TGF- α shedding responses (Data S1, $G\alpha_q$ (ΔC) conditions). Small interfering RNA (siRNA)-mediated knockdown experiments in the parental HEK293 cells confirmed involvement of $G_{q/11}$ and $G_{12/13}$ in the TGF- α shedding response (Figure S1). Thus, this is clear evidence that the TGF- α shedding assay selectively measures $G_{q/11}$ and/or $G_{12/13}$ signaling.

To exclude the possibility that the blunted AP-TGF- α release signal was caused by loss of GPCR expression, we compared surface expression levels of epitope-tagged GPCRs among parental, ΔG_q , ΔG_{12} , and $G_q/\Delta G_{12}$ cells using a flow cytometry. All tested GPCRs (FLAG-ADRB1, FLAG-HRH1, and FLAG-DRD1) were equally expressed in the parental as well as the G-protein-knockout (KO) cells (Figure S2A).

Chimeric G-Protein-Based Signaling Assay

We exploited the above assay system, and the previously identified importance of the $G\alpha$ subunit C-terminus, to develop the TGF- α shedding assay to assess binding of G proteins to any GPCR of interest. Specifically, we constructed chimeric $G\alpha$ subunits where the native 6-amino acid C-termini of members from the $G\alpha_{q/11}$ and the $G\alpha_{12/13}$ families were substituted with those from other human $G\alpha$ subunits (Figures 1C and S2A–S2C) and expressed them together with a test GPCR in the signaling-silenced $\Delta G_q/\Delta G_{12}$ cells (Figure 1C). The resulting downstream

coupling is scored as logarithmic values of relative intrinsic activity (RAI), which is defined as a relative E_{max}/EC_{50} value normalized by the highest value. Symbol size is proportional to E_{max} , which reflects fitting quality. During data processing, a concentration-response curve that failed to converge or had an E_{max} value of less than 3% AP-TGF- α release, or a RAI value of less than 0.01 was defined as LogRAI value of -2 . Data for the concentration-response curves are from a representative experiment (mean \pm SD of triplicate measurements). Each LogRAI plot denotes a single experiment and bars and error bars represent mean and SEM, respectively ($n = 4$).

See also Figures S1 and S2 and Data S1, S2, and S3.

signals measured by the TGF- α shedding assay should thus reflect the true binding events between any GPCR and its G-protein counterparts (Figure 1D).

We tested a series of chimeric G α subunits for their ability to induce the TGF- α shedding response. Specifically, we constructed chimeric G α subunits with the same C-terminal tail, but a different backbone (Figures S2D and S2E), using members of the G $_{q/11}$ (G α_q , G α_{11} , G α_{14} , and G α_{16} subunits) and the G $_{12/13}$ (G α_{12} and G α_{13} subunits) subfamilies. We expressed each chimeric G α subunit (C-terminal G α_{11} or G α_s chimeras) together with a test GPCR (G $_{i/o}$ -coupled DRD2 or G $_s$ -coupled PTGER2, respectively) and stimulated the cells with an agonist. We found that the G α_q backbone was the most efficacious in inducing TGF- α shedding response (%AP-TGF- α release response) for both receptors (Figures S2D and S2E); we thus chose this backbone for all subsequent experiments.

We generated chimeric G α_q subunits for each of the 11 unique C-terminal hexapeptides, which cover all of the 16 human G α subunits (Figures 1C and S2A–S2C; the C-terminal 6-amino acids are identical for G α_{11} , G α_{12} , G α_{11} , G α_{12} , and G α_{13} ; and for G α_q and G α_{11}), and one negative control lacking the tail (G α_q Δ C). The 11 G α_q chimeras were equally expressed in Δ G $_{q/12}$ cells (Figure S2F). Transfected cells were harvested and seeded in a 96-well plate and stimulated with or without titrated concentrations of a GPCR ligand (typically, 12 points in total). AP-TGF- α release signals over titrated concentrations were fitted with a sigmoidal concentration-response curve, from which EC $_{50}$ and E_{max} (an amplitude of ligand-induced response) values were obtained. For each chimeric G α condition, an E_{max}/EC_{50} value was normalized by the maximum E_{max}/EC_{50} value among the 11 G α chimeras (e.g., G α_q C terminus for TBXA2R; Figure 1D). This gave a relative, dimensionless E_{max}/EC_{50} value (relative intrinsic activity, RAI [Ehlert et al., 1999]), which was then base-10 log-transformed (LogRAI) and used as G-protein-coupling indices. With our pre-determined threshold criteria (STAR Methods), LogRAI ranged from -2 to 0 (100-fold in linear range). The assay produced robust, reproducible results as evidenced by well-clustered plots across independent experiments (Figure 1D; Data S1). By using a similar approach (restoration of a chimeric G α subunit in G-protein-KO cells), we performed a G $_s$ -based cAMP assay in G $_s$ -deficient (Δ G $_s$) HEK293 cells and confirmed that LogRAI values obtained from the G α_s backbone and the G α_q backbone were well correlated in prostanoid receptors (Figures S2I–S2K; Data S2; $r^2 = 0.74 \pm 0.16$, $n = 7$), which show distinct G-protein-coupling profiles (Sugimoto and Narumiya, 2007; Woodward et al., 2011).

As above, we measured cell-surface expression of GPCRs by flow cytometry to exclude GPCR expression level effects (Figure S1F). Except for a modest increase in the conditions with the chimeric G α_{13} co-expression, expression levels of N-terminal FLAG epitope-tagged GPCRs (FLAG-ADRB1 and FLAG-HRH1) were almost equal among cells co-expressing any of the G α chimeras.

The NanoBiT-G-Protein Dissociation Assay

To complement the chimeric G-protein-based TGF- α shedding assay, we established an additional assay in which dissociation of the G α subunit from the G $\beta\gamma$ subunits, a critical process of

G-protein activation, is measured via a luciferase complementation system. Bioluminescence resonance energy transfer (BRET) between a *Renilla* luciferase-inserted G α subunit and a GFP10-fused G β or G γ subunit was previously developed to measure G α -G $\beta\gamma$ dissociation (Galés et al., 2005). Here, we replaced the BRET pair with a split luciferase (NanoLuc Binary Technology; NanoBiT) (Dixon et al., 2016). Specifically, we inserted a large fragment (LgBiT) of the NanoBiT into the helical domain (between the α A and the α B helices) of a G α subunit (G α -Lg) and fused a small fragment (SmBiT) to the N-terminus of G β or G γ subunit (Sm-G β or Sm-G γ) (Data S3). We confirmed that G α_s -Lg retained a G $_s$ signaling function by measuring adenylyl cyclase-activating activity in Δ G $_s$ cells upon G $_s$ -coupled-receptor stimulation (Figures S2G and S2H). We generated a series of G α -Lg, Sm-G β , and Sm-G γ subunits (Data S3) and optimized a combination (STAR Methods). When expressed together in cells, these constructs form a heteromer possessing a complemented enzyme, whose luciferase activity is measurable upon loading with coelenterazine (CTZ), a substrate of the luciferase (Figure S3A). GPCR ligand stimulation triggers dissociation of G α -Lg from Sm-G β /G γ , making the real-time dissociation response detectable (Figure S3B). The NanoBiT-G-protein assay demonstrated highly reproducible dissociation signals across independent experiments and was minimally affected by preincubation time with CTZ (Figure S3F). Comparison of NanoBiT-G-protein dissociation signals with the chimeric G-protein-based assay for eight prostanoid receptors showed a moderately strong correlation ($r^2 \geq 0.5$) across all of the four G-protein subfamilies (Figures S3F and S3G).

Ligand Biased G-Protein Signaling

Since the chimeric G-protein-based assay recognize a ligand-activated conformation of a GPCR, we assessed whether it could also detect ligand bias among different G-protein subfamilies. An angiotensin II (Ang II) analog, [Sar¹, Ile^{4,8}]-Angiotensin II (SII), was shown to induce G $_{i/o}$ over G $_{q/11}$ as compared with Ang II in cells expressing AGTR1 (Saulière et al., 2012). We performed the assay using Ang II and SII (Figure S1G) and calculated coupling scores for Ang II-induced LogRAI and SII-induced LogRAI (Figure S1H). If SII behaves as a balanced agonist across G proteins, LogRAI plots obtained from Ang II and SII would be linearly aligned. Our results recapitulated the G $_i$ -bias of SII (Saulière et al., 2012) and further showed that SII was biased toward G $_{12}$ over G $_q$ as compared with the reference ligand (Ang II). These findings were backed up by the NanoBiT-G-protein assay (Figure S1I).

Hundreds of Known and New Couplings

Using the chimeric G-protein-based assay, we profiled coupling across 148 human GPCRs (Figure 2; Tables S1A and S1B), which represent $\sim 80\%$ of liganded class A GPCRs. Whenever possible, we used endogenous ligands; when ligands were unstable (e.g., thromboxane A2 for TBXA2R) and/or endogenous ligands were not yet identified, we chose available synthetic ligands (U-46619 for TBXA2R and MDL29951 for GPR17).

We compared coupling data from the chimeric G-protein-based assay with that of GtoPdb. For each of the four G-protein subfamilies, we defined positive coupling if any member of the subfamily scored LogRAI ≥ -1 and negative coupling if all of




Figure 2. Signatures of G-Protein Coupling Determined by the Chimeric G-Protein-Based Assay

Heatmap of the LogRAi values for the 148 receptors of the chimeric G-protein-based assay. Cell colors range from blue (LogRAi = -2) to red (LogRAi = 0). Receptors and G proteins are rearranged according to the dendrogram of the full linkage clustering of the distance matrix calculated from the coupling profiles. Receptor gene symbols are colored according to family membership as reported in GtoPdb. See also [Figure S4](#) and [Tables S1](#) and [S2](#).

the members scored LogRAi < -1 ([Figures 3A](#) and [3B](#)). Receiver operating curve (ROC) analysis gives area under the curve (AUC) = 0.78 ([Figure S4A](#)) when considering high-confidence known coupling data and suggested a threshold of LogRAi \geq -1.0 (optimizing true positive rate [TPR] while minimizing false positive rate [FPR]; [STAR Methods](#); [Table S1D](#)) for defining true couplings. The assay also showed other broad similarities to GtoPdb, including $G_{i/o}$ being the most common, and $G_{12/13}$ the least ([Figures 3C](#) and [3D](#)). We also recapitulated that the majority of receptors coupled to only one G-protein, which does not change greatly with altered LogRAi thresholds ([Figure S4C](#)), though as expected there is greater coupling promiscuity at lower values. In addition, both GtoPdb and our data (at various LogRAi stringencies) suggest $G_{i/o}$ subunits to be the most specific, always displaying the highest fraction of exclusively bound receptors, with $G_{12/13}$ being the most promiscuous ([Figures 3C](#), [S4B](#), and [S4D](#)). A total of 39 promiscuous receptors are reported to couple to members of all four G-protein families ([Figure 4C](#)); however, promiscuity decreases as a function of the LogRAi threshold ([Figures S4E](#) and [S4F](#)). Overall, the dataset shows an excellent agreement with known couplings ([Figure 3D](#)), with more than 88% of reported couplings reproduced for three classes (i.e., $G_{i/o}$, $G_{q/11}$, and $G_{12/13}$).

As seen in previous studies ([Flock et al., 2017](#); [Horn et al., 2000](#); [Rodriguez et al., 2010](#)), we found no correlation between sequence and coupling similarities, either performing pairwise

comparisons on the whole set or intra-family ([Figures 2](#) and [S4G](#)). Moreover, both extremes are evident: receptor pairs with low sequence similarity can have similar couplings and close homologs from the same family can show large differences (see prostanoid receptors; [Figures 3A](#) and [3B](#)). Our exploration of 11 distinct G proteins also reveals key differences among G-protein subfamilies in terms of their coupling preferences, which essentially reflects sequence similarity of the last 6 C-terminal amino acids ([Figures 2](#) and [S2C](#)). For instance, several receptor families show overall coupling preferences for specific classes, like Opioid and Dopamine receptors for $G_{i/o}$, or Prostanoid and Adrenoceptors for G_s ; in contrast others show more coupling promiscuity, like Endothelin, Ghrelin, and Proteinase-activated receptors ([Figure 2](#); [Data S1](#)). The great utility of our dataset is immediately clear. There are entire groups of poorly annotated (in GtoPdb) receptors that are well represented in our dataset, including ten GPCRs ([Table S1C](#)) such as protease-activated receptors and P2Y receptors (P2RY10 and P2RY12), where the latter is a major target of antiplatelet agents, with roles in platelet aggregation ([Dorsam and Kunapuli, 2004](#)) and bleeding disorders ([Patel et al., 2014](#)). While P2RY10 displays specificity for both the $G_{i/o}$ and the $G_{12/13}$ subfamily members, we find P2RY12 to be specific for $G_{i/o}$ ([Tables S1B](#) and [S1C](#)). Elsewhere, GPR132, recently emerged as a mediator of breast carcinoma metastasis ([Chen et al., 2017](#)), shows a previously unreported coupling promiscuity ([Table S1B](#)).




Figure 3. Comparison of G-Protein Coupling between the Chimeric G-Protein-Based Assay and GtoPdb, and Validation of G_{12/13} Signaling for the Newly Characterized GPCRs

(A) Classification of the LogRAi scores and its comparison with GtoPdb. An example heatmap of LogRAi scores for the eight prostanoid receptors is shown, with a LogRAi cutoff of -1 to binary-classify the data into coupled (red-to-white; Y) or uncoupled (blue-to-white) classes. G-protein coupling from GtoPdb (subfamily levels) is overlaid.

(B) Combined binary coupling/non-coupling data for each of the four G-protein subfamilies.

(C) Venn diagrams with the numbers of receptors coupled to each G-protein subfamily in the chimeric G-protein-based assay (LogRAi ≥ -1).

(D) Venn diagrams of receptor couplings to the four G-protein families according to the chimeric G-protein-based assay (LogRAi ≥ -1) and GtoPdb.

(E) GPCRs that were identified as being coupled with G_{12/13} by the chimeric G-protein-based assay were examined for their ability to engage and activate native, endogenous G_{12/13} in HEK293 cells. A test GPCR was expressed in the parental, ΔG_q , ΔG_{12} , and $\Delta G_q/\Delta G_{12}$ cells along with the AP-TGF- α reporter construct, but not with a chimeric G α subunit, and its ligand-induced response was assessed. Note that, in all of the tested GPCRs, AP-TGF- α release response occurred in ΔG_q cells but was completely silenced in $\Delta G_q/\Delta G_{12}$ cells, showing induction of G_{12/13} signaling. Symbols and error bars represent mean and SEM, respectively, of 3–6 independent experiments with each performed in triplicate.

See also Figures S4 and S5 and Tables S1.

identified G_{12/13} couplings indeed reflect capability of endogenous G_{12/13} activation, and not artifacts of chimeric G α subunit overexpression, we assessed TGF- α shedding responses in ΔG_q cells (Figures 1A and 1B). We tested eight GPCRs (AGTR1, CNR1, EDNRA, F2RL2, PTGER1, PTGFR, TACR1, and TBXA2R), in which G_{12/13} coupling was not registered in GtoPdb but was detected by the chimeric G-protein-based assay. We found that all of them induced TGF- α shedding responses in ΔG_q cells and that the signals were completely silenced in $\Delta G_q/\Delta G_{12}$ cells (Figure 3E). To assess a more proximal signaling event to G_{12/13} activation, we generated a NanoBit-RhoA sensor (Figures S5A–S5C; STAR Methods) and found that all of tested

In general, more than half of the couplings detected (160/292, 55%) have not previously been reported (Figure 3D). As expected, the biggest proportion of new couplings are G_{12/13} where our data make up 57% of all known couplings of this type (15 out of 26 reported G_{12/13} couplings in GtoPdb, which also considers non-class A GPCRs). To validate that newly

GPCRs that were newly identified as G_{12/13}-coupled receptors (Figure 3E), when overexpressed in HEK293 cells, induced RhoA activation (Figure S5D). We also found that thrombin activated RhoA, presumably by activating its receptors (F2L, F2RL2, and/or F2RL3) that were endogenously expressed in PC-3 and MDA-MB-231 cells (Figure S5E). In HN12 cells and




Figure 4. Development of G-Protein-Coupling Predictor

(A) Workflow of the procedure: features are extracted from sub-alignments of coupled and uncoupled receptors to a particular G protein; features are used to generate a training matrix, which is employed to train a logistic regression model through a 5-fold cross-validation procedure.

(B) The final model is tested on reported couplings not previously seen during training and compared to PredCouple.

(C) Highly confident predicted couplings (coupling probability >0.9) for 61 class A GPCRs lacking information about transduction from both GtoPdb or the chimeric G-protein-based TGF- α shedding assay (black) versus receptors with experimental coupling information (gray).

See also Figure S7, Table S2, and Data S4.

Cal27 cells, a RhoA pull-down assay showed that CP-55940 (CP; likely via CNR1, but not CNR2; Figure S5E) and Ang II induced activation of RhoA (Figure S5F). Together, these data demonstrate that the G_{12/13}-coupled receptors identified by the chimeric G-protein-based assay induce RhoA activation at both overexpression and endogenous expression levels.

To test whether apparent unchanged TGF- α shedding responses in Δ G₁₂ cells as compared with those in the parental cells (AGTR1 and EDNRA) arose from enhanced or compensated G_{q/11} signaling in Δ G₁₂ cells, we analyzed G-protein expressions and performed a Ca²⁺ mobilization assay and a NanoBiT-IP₃ assay (Figure S6; STAR Methods), both of which selectively measure G_{q/11} signaling (Figures S6C and S6F). Expression levels of G α_q and G α_{11} were unchanged in Δ G₁₂ cells, nor was that of G α_{13} in Δ G_q cells (Figure S6A). Both Ca²⁺ and IP₃ responses in Δ G₁₂ cells were comparable to those in the parental cells (Figures S6C and S6G). Thus, TGF- α shedding responses in Δ G₁₂ and Δ G_q cells are an accurate reflection of G_{q/11} and G_{12/13} signaling.

Sequence Features Indicative of Coupling Specificity

We used a statistical model to identify sequence features associated with each of the eleven couplings determined above (all details given in STAR Methods). Briefly, we used sequence alignments for each coupling group to define residues and more general compositional features (e.g., C-terminal or IC3 length, charge distributions, etc.) found to be statistically associated to coupling for each G protein. These were used to train and test a machine learning (Logistic regression) predictor (Figure 4A) and identify the features most predictive for each G protein.

Our predictor performs better than another available approach (PredCouple) (Sgourakis et al., 2005a) in predicting known couplings not used during training for all coupling groups, but particularly for G_{12/13}, which is expected since few data were available to train such predictors previously (Figure 4B; Tables S2A and S2C). Note that same predictor trained only with known couplings from GtoPdb (Harding et al., 2018) performed worse (Figure 4B; Table S2D) as might be expected. Using a stricter LogRAi cutoff to define coupling groups led to a general decrease in performance during the testing phase, except for

$G_{i/o}$ (Figures S7A–S7C). The poorer performance, for the G_s subfamily, which also shows a poorer overlap between the chimeric G-protein-based assay and GtoPdb (Figure 3D), is probably a consequence of the fact that the chimeric system does not capture all sequence determinants emerging for G_s coupling (Carpenter et al., 2016; García-Nafria et al., 2018a; Rasmussen et al., 2011). Nevertheless, this tool can be exploited to illuminate the transduction mechanisms of less-characterized receptors. Indeed, for the 61 receptors (21% of 286 class A GPCRs) lacking coupling information from either GtoPdb or the chimeric G-protein-based assay (Table S1C), we predict a prevalence of G_s followed by $G_{q/11}$ and $G_{12/13}$ couplings, the latter contrasting with the smallest fraction among experimental couplings (Figure 4C; Table S2E). For example, P2RY8 is readily predicted to be coupled to $G_{12/13}$ (Table S2E), being consistent with a report of mutual exclusive mutations in lymphomas between the *P2RY8* and the *GNA13* genes, which implies a putative functional link (Muppidi et al., 2014).

The model identified different combinations of sequence features important for each coupling group (Figure 5). After training, different weights are assigned to each feature in the logistic function to achieve optimal prediction performances, thus highlighting the most relevant determinants for each coupling (Figure 5; STAR Methods). Significant features are more abundant at the cytosolic side of the receptor (Figures 6A and 6B; Table S3A) including many at the known G-protein-binding interface (e.g., transmembrane helix 3 [TM3], TM5, TM6, and intracellular loop 3 [ICL3]), but also within the core of the structure, mainly contributing to a contact network (Venkatakrisnan et al., 2016) and could thus mediate specific conformational differences required to accommodate a particular G-protein (Figure 6).

Surprisingly, only a few significant positions (12 of 51 or 23%) overlap with residues lying directly at known GPCR/G-protein interfaces (Figures 5A and 6C; Table S3A). These include ICL3, TM5, and TM6 positions associated with $G_{i/o}$ missing from G_s (e.g., 5.61, ICL3:174, 191–194) that are likely responsible for specificity (Draper-Joyce et al., 2018; García-Nafria et al., 2018b). Several other positions (11, 21%) are immediately adjacent to direct contacts, suggesting they could nevertheless affect these interfaces (Table S3B). This is logical as some of the contacting positions are typically highly conserved across GPCRs (e.g., the DRY or NPxxY motifs). Overall, the majority (or 90%) of significant positions within the 7TM bundle mediate intra- or inter-protein contacts with either G proteins or ligands. The majority of significant positions (29, 57% of the total) appear to mediate active-like state specific intramolecular contacts, which we uncovered by comparing functional state specific contact networks (i.e., active-like and inactive-like) from three-dimensional (3D) structures (Figures 6C–6E, Tables S3A and S3B; STAR Methods). Helices TM3, TM5, and TM6 undergo major rewiring of their intramolecular contacts upon receptor activation and display the highest content of significant coupling features in the active-like network (Figure 6D). This further stresses their role as master regulators of receptor activation and G-protein recognition (Koehl et al., 2018). Residues previously described as universal mediators of receptor activation (Venkatakrisnan et al., 2016) participate to this network as either endpoints (6.37) or mediators (3.46 and

7.53) of the shortest paths linking the ligand and G-protein-binding pockets (Figures S7D and S7E; Table S3B; STAR Methods).

Several other features lie within regions outside the 7TM bundle, particularly in the ICL3 or C-terminal regions (Figures 5C, 6A, and 6B), that are not usually visible in experimental structures (with the exception of ICL3 in some $G_{i/o}$ complexes), but which nevertheless play critical roles in signaling (Venkatakrisnan et al., 2014). There is broadly an equal contribution of positions from within or outside of the 7TM bundle across all families, with a greater prevalence of the outside positions for the $G_{12/13}$ subfamily (Figures 6A and 6B).

Data-Driven Design of a G_{12} -Specific DREADD

The prominent roles for ICL3, and to a lesser extent the C terminus, for $G_{12/13}$ -coupled receptors, where length and electrostatic charge are predicted to be important for coupling (Figures 5B and 5C), together with the lack of structure and tools for probing $G_{12/13}$ signaling prompted us to develop a new chemogenetic receptor for studying $G_{12/13}$ coupling. DREADDs are engineered receptors that permit spatial and temporal control of G-protein signaling *in vivo*, being thus of great use in studying and manipulating signaling (Urban and Roth, 2015; Wess et al., 2013). DREADDs derived from the muscarinic acetylcholine (ACh) receptors are widely used in combination with clozapine N-oxide (CNO), a synthetic, biologically inert ligand. To date, DREADDs coupling to G_s , $G_{i/o}$, and $G_{q/11}$ (M3D- G_s , M4D, and M3D, respectively) have been developed (Armbruster et al., 2007; Guettier et al., 2009), but there is not yet a $G_{12/13}$ -coupled DREADD available, which we sought to design using our predictor.

The design of M3D- G_s involved a strategy of substituting both ICL2 and ICL3 of the $G_{q/11}$ -coupled M3D with those of G_s -coupled $\beta 1AR$ (Guettier et al., 2009). In our analysis, a major feature contributing to $G_{12/13}$ coupling was ICL3, followed by the C-terminal tail (Figures 5C and 6B). We thus explored whether these features would be sufficient to induce such signaling in M3D. We first predicted the probability of G_{12} coupling for M3D chimeras containing ICL3 swapped from all other GPCRs alone or in combination with C-terminal stretches (Figures 7 and S7F). Among all possible GPCR constructs (144 ICL3-swapped chimeras and 144 dual ICL3/C terminus chimeras), we selected the top 10 predictions of each chimera type (13 GPCRs in total by excluding overlaps and selecting representative constructs when multiple members from one GPCR were predicted) leading to 26 constructs (Figure 7A). We functionally screened G_{12} -coupling activity of the M3D-based chimeras using two assays. In the first, the chimera construct was expressed together with the AP-TGF- α reporter in ΔG_q cells, in which G_{12} signaling is selectively detectable (Figure 3E). We measured TGF- α shedding response upon CNO or ACh stimulation. Among the 26 constructs screened, chimeras with the GPR183-derived ICL3 substitution (M3D-GPR183/ICL3) and the GPR132-derived ICL3 substitution (M3D-GPR132/ICL3) showed significant G_{12} signaling (*p* values <0.05) (Figure 7B). ACh did not induce detectable G_{12} signaling in any of the tested constructs (Figure 7B). In the second assay, the chimera construct was expressed together with NanoBiT- G_{12} (in combination with Sm- $G_{\gamma 11}$) in




Figure 5. Featured Residues in GPCRs Involved in G-Protein-Coupling Selectivity

(A) Comparison of significant coupling features weights for the 11 G proteins (bottom), interface contacts of 6 available GPCR-G-protein complexes (central) and 7TM domain position conservation (top). On the bottom panel are all the features (columns) that are found to be statistically significant ($p < 0.05$) for at least one coupling group (rows). Each cell is colored based on coefficient of the given feature in the decision function of the corresponding coupling group (i.e., weight), with negative and positive values colored red and green respectively. Coupling features at 7TM domain with significantly different amino acid distributions are characterized by two values, representing the weights of the bitscores obtained from the coupled (top sub-cell) and not coupled (bottom sub-cell) hidden Markov modes (HMMs) for each G protein. Insertions (i.e., positions present only in the coupled subset) or deletions (i.e., positions present only in the uncoupled subset) are indicated with a gray “+” and “-.” Black/gray boxes in the center show contacts mediated by the last 6 amino acids of C-terminal helix 5 of G α subunits (black) and contacts mediated by the other positions of G α subunits (gray). Top bars shows conservation profiles for PFAM 7tm_1 positions obtained by calculating the information content from HMM positions bit scores (Wheeler et al., 2014).

(B) 2D cartoons of the 7TM topology indicating the regions contributing to the features. ICL, intracellular loop; ECL, extracellular loop; TM, transmembrane helix.

(C) Significant coupling feature weights for the 11 G proteins (same color codes as in A) of extra-7TM features of ICL3 and C terminus, including length and amino-acid composition.

See also Figures S4 and S7 and Table S3.

the parental HEK293 cells and stimulated with CNO. As a counter experiment, we used NanoBiT-G_o since our preliminary experiment indicated a minor coupling of some chimeras to G_o. NanoBiT-G₁₂ screening identified four constructs (M3D-GPR183/ICL3, M3D-GPR132/ICL3, M3D-P2RY10/ICL3, and M3D-NMBR/ICL3) that were significantly ($p < 0.05$) coupled to G₁₂ and not to G_o (Figure 7C). One construct (M3D-LTB4R2/ICL3) induced both G₁₂ and G_o coupling with a higher G_o dissociation signal. In both assays, dually swapped ICL3/C-terminus

chimeras showed negligible G₁₂ signaling, which was in part attributable to lower surface expression of these constructs (Figure 7D).

We then evaluated selectivity of G-protein coupling for the two candidate constructs using the NanoBiT-G-protein dissociation assay with titrated CNO concentrations. As controls, we compared with previously established muscarinic DREADDs (M3D, M4D, and M3D-G_s) (Armbruster et al., 2007; Guettier et al., 2009). We tested representative NanoBiT-G-proteins (G_s,




Figure 6. Functional Analysis of Residues Linked to Coupling Selectivity

(A) Upper panel: distribution of coupling feature fractions for extra- and intra-7TM portions. The formers comprise the 7TM helical bundle only, while the latter include the N and C termini, extra- and intra-cellular loops (ECLs and ICLs); lower panel: distribution of the coupling feature fractions within transmembrane sectors (i.e., EC, extracellular; TM, transmembrane; IC, intracellular). Extra- and intra-cellular portions are defined by ECL and ICL regions plus 5 helical positions preceding and following them (see Tables S3A and S3C).

(B) Distribution of the fractions of coupling significant features outside of the 7TM bundle.

(C) Distribution of coupling feature fractions (relative to the total number of positions of the same class) within functional sites (i.e., mediating either ligand/G-protein binding or intra-molecular contacts).

(D) Graph representing intra-molecular contacts within 7TM helices. Each helix is represented by a node, whose diameter is proportional to the number of helix positions mediating contacts in the contact network derived from active-like structures and whose color (red scale) is proportional to the number of significant coupling features present in the corresponding region. Edges represent contacts between 7TM helices, where width is proportional to the number of contacts in the active-like contact network, while color scale (gray) is proportional to the similarity degree (calculated as a Jaccard index) between contacts mediated in the active- and inactive-like contact networks.

(E) 3D cartoons of the *ADRB2*-*GNAS* complex (PDB: 3SN6) (Rasmussen et al., 2011) with side chains of coupling features at G-protein-binding sites depicted as red surfaces. Network drawings

were done through Cytoscape (<https://cytoscape.org/>). A representative coupling feature at intra-molecular contacts sites (i.e., position 3.40) is depicted as a red sphere mediating one of the shortest paths linking the ligand and G-protein-binding pockets (wheat sticks and spheres). The ligand and *GNAS* (G_{α_s}) are depicted as cyan and pale-yellow surfaces, respectively. 3D drawing was generated through pymol (<https://pymol.org/>). See also Figure S7 and Tables S2 and S3.







G_o , G_q , and G_{12}) from the four subfamilies. The NanoBIT-G-protein assay correctly measured primary coupling of the three established DREADDs (M3D, M4D, and M3D- G_s for G_q , G_o , and G_s , respectively; Figure 7E). M3D-GPR183/ICL3 and M3D-GPR132/ICL3 constructs showed robust G_{12} dissociation signals, while dissociations of the other G-proteins were much lower than those of G_{12} . None of the DREADD constructs induced significant NanoBIT- G_{13} dissociation (Figure 7E). The EC_{50} values of CNO for each primary-coupling G-protein were in a submicromolar range (0.1–1 μ M) for all of the DREADDs (Figure 7F). Thus, the constructs M3D-GPR183/ICL3 and M3D-GPR132/ICL3 are new G_{12} -selective DREADDs.

DISCUSSION

The extensive dataset provided here greatly expands known GPCR/G-protein couplings and provides better resolution by considering all 11 specific human G proteins rather than subfamilies. The assays, the resource, and accompanying predictor

(available at <http://gpcr.russelllab.org/>) can be used for a host of biological and pharmaceutical applications. For example, the TGF- α shedding assay, applied to AGTR1, demonstrates the promise to develop sub-G-protein-biased ligands (i.e., discriminating one G-protein signaling from another), which have recently attracted attention because of their potential for therapeutic-signal-targeted medicine with reduced on-target side effects (Violin et al., 2014). Most importantly, the extensive dataset provides the first coupling information for many receptors (e.g., protease-activated or P2Y receptors), shows differences in G proteins in the same family (e.g., prostanoïd receptors), and, in particular, identifies dozens of receptors coupled to the previously understudied $G_{12/13}$ (Rho signaling).

The $G_{12/13}$ subfamily remains challenging to study owing to lack of well-established methods for assessing signaling. The TGF- α shedding assay combined with ΔG_q cells is an excellent platform for selective measurement of $G_{12/13}$ signaling with high robustness and throughput and in the future will enable precise characterization of receptors and their ligands. Other assays



(legend on next page)

developed in this study (the chimeric G-protein-based TGF- α shedding assay, the NanoBiT-G-protein dissociation assay and the NanoBiT-RhoA sensor) will also be useful for cross-validating results. G_{12/13} signaling is also implicated in immune processes and various diseases (Herroeder et al., 2009; Suzuki et al., 2009), including receptors S1PR2 and P2RY8 in B cell lymphoma (Muppidi et al., 2014; O'Hayre et al., 2016). Agonists for G_{12/13}-coupled receptors in lymphocytes can attenuate immune responses and antagonists could potentially boost them, both of which offer attractive possibilities for future therapies. A list of the expanded members of G_{12/13}-coupled receptors identified here will provide a basis for such drug development. Indeed, some of our newly identified G_{12/13}-coupled GPCRs (e.g., CNR1, FFAR1, GHSR, GPR35, HRH2, HTR2C) are already targets for agonists approved as therapeutics (Hauser et al., 2017), suggesting additional possibilities for drug repurposing. Transgenic mice expressing our new G₁₂-coupled DREADD could help to explore G₁₂ signaling and ultimately develop such therapies.

Integrating this large GPCR/G-protein dataset with information about protein sequence and structure has identified numerous insights into how receptors selectively interact with G proteins. Several recent structures have provided insights into the complex landscape governing GPCR coupling specificity, which is complicated by multiple factors including conformational plasticity, kinetics, ligand biasing, and G-protein pre-association (Capper and Wacker, 2018). While previous efforts successfully identified sequence and structural features that determine coupling selectivity in *G proteins* (i.e., the barcode) (Flock et al., 2017), a systematic identification of *receptor* determinants is still lacking (Flock et al., 2017; Horn et al., 2000; Wong, 2003). Our analysis identifies several features that agree with what is already known. Generally, TM3, TM5, and TM6 have the greatest number of predicted coupling features, which is partly backed up by recent biophysical (Van Eps et al., 2018) and structural studies (Carpenter et al., 2016; Draper-Joyce et al., 2018; García-Nafria et al., 2018a, 2018b; Kang et al., 2018; Koehl et al., 2018; Rasmussen et al., 2011), suggesting the importance of ICL2, TM5, ICL3, and TM6 in determining complementarity to the G-protein barcode (Flock et al., 2017).

One potential issue with the results presented here is the use of chimeric G α subunits, where only the 6 C-terminal amino acids

are used to assess ligand-induced GPCR activation. This necessarily misses contributions of the remaining (backbone) region of the G α subunits, which are known to contribute to GPCR interaction (Carpenter et al., 2016; Draper-Joyce et al., 2018; García-Nafria et al., 2018a; Kang et al., 2018; Koehl et al., 2018; Rasmussen et al., 2011). However, the good agreement with known couplings (Figure 3D) suggests that these effects are not predominating. Moreover, relative contributions (or synergistic effects) of C terminus and backbone to coupling selectivity seem to differ among GPCRs (Figure 4A). Ultimately, an extensive G-protein-coupling dataset considering native α subunit sequences will naturally provide a more complete view of coupling determinants.

One would expect naively that coupling determinants would only lie at the interface between G proteins and receptors and that a few simple sequence changes would account for selectivity. Decades of sequence gazing have failed to find such simple explanations as have long been seen, for example, in enzyme/substrate specificity (Hannenhalli and Russell, 2000). Recent receptor/G-protein complexes suggest that additional features outside the interface, such as an internal network of polar contacts, induce a greater rigidity of TM6 and lead to a preference of G_{i/o} over G_s (Kang et al., 2018; Koehl et al., 2018). Many of our predicted sequence features away from the interface indeed participate in intra-molecular contact networks linking ligand and G-protein-binding sites, which broadly agrees with this notion and is supported by other contact-network analyses (Angelova et al., 2011; Venkatakrishnan et al., 2013, 2016). We speculate that these features allow allosteric and dynamic control of a G-protein-binding interface of GPCRs possibly by stabilizing a specific intermediate state of a receptor/G-protein complex. We also find a general tendency for TM5, ICL3, and TM6 insertions in G_{i/o}-coupled and deletions in G_s-coupled receptors, which broadly agrees with the notion that the bulkier side-chains of the G_s C-terminus can only be accommodated by larger and more flexible crevices found in G_s-specific receptors (Figure 7A) (García-Nafria et al., 2018a; Kang et al., 2018).

We predicted many G-protein-coupling features to lie outside of the 7TM bundle. For example, ICL3 contains features for G_{12/13} coupling, the importance of which is verified by the successful generation of ICL3-swapped DREADDs. G_{12/13} is

Figure 7. Generation of G₁₂-Coupled Designer GPCRs

(A) Scheme of generating and assessing ICL3 or ICL3/C terminus-swapped constructs from G_{q/11}-coupled M3D. Based on the predictor scoring of 288 constructs from 144 GPCRs (Figure S7F), we selected 13 GPCRs and made 26 constructs.

(B–D) Functional screening of M3D-derived chimeric constructs. G₁₂ signaling of the constructs assessed by the TGF- α shedding assay in the Δ G_q cells treated with 10 μ M clozapine N-oxide (CNO) or 10 μ M acetylcholine (ACh) (B). Activation of G₁₂ and G_o was measured by the NanoBiT-G-protein dissociation assay with 10 μ M CNO (C). G α_{12} -Lg or G α_o -Lg was co-expressed with Sm-G γ t1 (Data S3). Changes in decreased luminescent signals are inversely plotted in the y axis. (C). Surface expression of the M3D-derived chimeric constructs was assessed by a flow cytometry using an anti-FLAG epitope-antibody, followed by a fluorescently labeled secondary antibody (D). Symbols and error bars represent mean and SEM, respectively, of 4–8 independent experiments with each performed in duplicate or triplicate. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$ (two-way ANOVA, followed by Sidak's multiple comparison tests).

(E) Lack of G₁₃ activation by the new DREADD constructs. Dissociation signals of the NanoBiT-G₁₃ protein were assessed by using 10 μ M CNO (M3D-GPR183/ICL3 and M3D-GPR132/ICL3) and 1 μ M U-46619 (TBXA2R). Symbols and error bars represent mean and SEM, respectively, of 3–11 independent experiments with each performed in duplicate.

(F) Concentration-response curves for G-protein activation by DREADD constructs. Previously established DREADDs (G_{q/11}-coupled M3D, G_{i/o}-coupled M4D, and G_s-coupled M3D-G_s) and the newly generated DREADDs (M3D-GPR183/ICL3 and M3D-GPR132/ICL3) were profiled for their G-protein coupling using representative members (G_s, G_{i1}, G_q, and G₁₃) of the 4 G-protein subfamilies. Symbols and error bars represent mean and SEM, respectively, of 3–12 independent experiments with each performed in duplicate. For each DREADD, parameters for the most efficaciously coupled G protein are shown in bottom of the panel. See also Figure S7.

the receptor class where we predict the smallest number of significant features overlapping with G-protein interface residues (Figures 5A and 6C) and the greatest fraction of features outside the 7TM bundle, particularly in ICL3 (Figure 6A). Since the ICL3 is typically disordered (i.e., lacks a pre-defined structure), it is possible that the fewer specific couplings observed for G_{12/13} receptors (Figure 3C) are a consequence of the lack of well-defined contact points in the receptor structure. Since (non G_{12/13}) G-protein/GPCR complex structures show limited but nevertheless G-protein-class-specific interactions between ICL3 and flanking amino acid residues (i.e., TM5/ICL3 for G_s and ICL3/TM6 for G_{i/o} complexes; Figure 5A), we speculate that G_{12/13} receptors might also engage in ICL3 and Helix 5 in the G α subunit-specific interactions that are likely different from G_s or G_{i/o} (or that an ICL3-Helix 5 interaction occurs during an intermediate state).

This study has demonstrated the power of integrating a new, powerful assay with systematic data analysis to provide new insights in molecular mechanism. With the extensive analysis, we devised both biological and computational tools that will advance understanding of how cells respond to extracellular signals. Integrating our resource with other datasets, such as genomic sequencing, transcriptomics, proteomics, metabolomics, and/or by considering other members of GPCRs mediated pathways, will provide new means to quantify downstream signaling in normal and pathological conditions and provide considerable possibilities for new therapies and personalized medicine.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- **KEY RESOURCES TABLE**
- **CONTACT FOR REAGENT AND RESOURCE SHARING**
- **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
 - Cells and transfection
- **METHOD DETAILS**
 - Plasmids
 - TGF- α shedding assay
 - Calculation of G-protein-coupling score
 - NanoBiT-G-protein dissociation assay
 - NanoBiT-RhoA assay
 - NanoBiT-IP₃ sensor assay
 - siRNA transfection
 - Quantitative real-time PCR analysis
 - Western blot
 - Flow cytometry
 - GloSensor cAMP assay
 - Active RhoA pulldown assay
 - Ca²⁺ mobilization assay
 - Comparison of data from the chimeric G-protein-based assay with known couplings
 - Sequence-based coupling-determinant features
 - G-protein-coupling predictor
 - Training and cross validation
 - Randomized training test

- Test set to benchmark predictor performance
- Functional classification of coupling features through 3D structure analysis
- Predictions of G₁₂-coupled DREADD chimeric sequences
- **QUANTIFICATION AND STATISTICAL ANALYSIS**
- **DATA AND SOFTWARE AVAILABILITY**
- **ADDITIONAL RESOURCES**

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.cell.2019.04.044>.

ACKNOWLEDGMENTS

We thank Frederick J. Ehlert at University of California, Irvine and Sudarshan Rajagopal at Duke University for helpful comments on analyses of the chimeric G-protein-based assay; Michel Bouvier at University of Montreal for discussion on G-protein profiling; members of the laboratory at Tohoku University for plasmid preparation; Ayumi Inoue at Tohoku University for flow cytometry analyses. This work was supported by JSPS KAKENHI grant 17K08264 (A.I.); Japan Science and Technology Agency (JST), PRESTO JPMJPR1331 (A.I.); and the PRIME JP17gm5910013 (A.I.) and the LEAP JP17gm0010004 (A.I. and J.A.) from the Japan Agency for Medical Research and Development (AMED). N.A. is supported by training grants from the National Institute of General Medical Sciences (NIGMS) T32GM007752 and the National Science Foundation (NSF) DGE-1650112. J.S.G. is supported by relevant grants from the National Cancer Institute (NCI) R33CA225291 and U54CA209891; and the National Institute of Dental and Craniofacial Research (NIDCR) U01DE028227. F.R., G.S., and R.B.R. are supported by the Cell Networks Excellence initiative of the Germany Research Foundation (DFG). R.B.R. is part of the DFG SFB/TPR186 Molecular Switches in the Spatio-Temporal Control of Cellular Signal Transmission and the BMBF German Network for Bioinformatics (de.NBI). F.R. was additionally supported by an Alexander Von Humboldt post-doctoral fellowship (F.R.) and a Michael J. Fox Foundation research grant (F.R. and R.B.R.).

AUTHOR CONTRIBUTIONS

Conceptualization, A.I. and F.R.; Methodology, A.I., F.R., G.S., J.A., and R.B.R.; Investigation, A.I., F.M.N.K., Y.S., S.I., T.K., F.R., A.U., Y.O., N.A., K.K., G.S., and J.S.G.; Writing, A.I., F.R., and R.B.R. with feedback from all of the coauthors; Funding Acquisition, A.I., J.A., F.R., and R.B.R.; Resources, A.I.; Supervision, A.I., F.R., and R.B.R.

DECLARATION OF INTERESTS

A.I., F.R., J.A., and R.B.R. have filed a patent application related to this work.

Received: October 25, 2018

Revised: January 28, 2019

Accepted: April 25, 2019

Published: May 30, 2019

REFERENCES

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment search tool. *J. Mol. Biol.* *215*, 403–410.
- Angelova, K., Felline, A., Lee, M., Patel, M., Puett, D., and Fanelli, F. (2011). Conserved amino acids participate in the structure networks deputed to intramolecular communication in the lutropin receptor. *Cell. Mol. Life Sci.* *68*, 1227–1239.
- Ambruster, B.N., Li, X., Pausch, M.H., Herlitze, S., and Roth, B.L. (2007). Evolving the lock to fit the key to create a family of G protein-coupled receptors

- potently activated by an inert ligand. *Proc. Natl. Acad. Sci. USA* **104**, 5163–5168.
- Ataei, F., Torkzadeh-Mahani, M., and Hosseinkhani, S. (2013). A novel luminescent biosensor for rapid monitoring of IP3 by split-luciferase complementary assay. *Biosens. Bioelectron.* **47**, 642–648.
- Capper, M.J., and Wacker, D. (2018). How the ubiquitous GPCR receptor family selectively activates signalling pathways. *Nature* **558**, 529–530.
- Carpenter, B., Nehmé, R., Warne, T., Leslie, A.G., and Tate, C.G. (2016). Structure of the adenosine A(2A) receptor bound to an engineered G protein. *Nature* **536**, 104–107.
- Chen, P., Zuo, H., Xiong, H., Kolar, M.J., Chu, Q., Saghatelian, A., Siegwart, D.J., and Wan, Y. (2017). Gpr132 sensing of lactate mediates tumor-macrophage interplay to promote breast cancer metastasis. *Proc. Natl. Acad. Sci. USA* **114**, 580–585.
- Conklin, B.R., Farfel, Z., Lustig, K.D., Julius, D., and Bourne, H.R. (1993). Substitution of three amino acids switches receptor specificity of Gq alpha to that of Gi alpha. *Nature* **363**, 274–276.
- Csardi, G., and Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal Complex Systems* **1695**. <http://igraph.org>.
- Denker, B.M., Schmidt, C.J., and Neer, E.J. (1992). Promotion of the GTP-liganded state of the Go alpha protein by deletion of the C terminus. *J. Biol. Chem.* **267**, 9998–10002.
- Devost, D., Sleno, R., Pétrin, D., Zhang, A., Shinjo, Y., Okde, R., Aoki, J., Inoue, A., and Hébert, T.E. (2017). Conformational Profiling of the AT1 Angiotensin II Receptor Reflects Biased Agonism, G Protein Coupling, and Cellular Context. *J. Biol. Chem.* **292**, 5443–5456.
- Dijkstra, E. (1959). A note on two problems in connexion with graphs. *Numer. Math.* **1**, 269–271.
- Dixon, A.S., Schwinn, M.K., Hall, M.P., Zimmerman, K., Otto, P., Lubben, T.H., Butler, B.L., Binkowski, B.F., Machleidt, T., Kirkland, T.A., et al. (2016). NanoLuc Complementation Reporter Optimized for Accurate Measurement of Protein Interactions in Cells. *ACS Chem. Biol.* **11**, 400–408.
- Dorsam, R.T., and Kunapuli, S.P. (2004). Central role of the P2Y12 receptor in platelet activation. *J. Clin. Invest.* **113**, 340–345.
- Dou, Y., Wang, J., Yang, J., and Zhang, C. (2012). L1pred: a sequence-based prediction tool for catalytic residues in enzymes with the L1-logreg classifier. *PLoS ONE* **7**, e35666.
- Draper-Joyce, C.J., Khoshouei, M., Thal, D.M., Liang, Y.L., Nguyen, A.T.N., Furness, S.G.B., Venugopal, H., Baltos, J.A., Plietzko, J.M., Danev, R., et al. (2018). Structure of the adenosine-bound human adenosine A₁ receptor-G_i complex. *Nature* **558**, 559–563.
- Eddy, S.R. (1998). Profile hidden Markov models. *Bioinformatics* **14**, 755–763.
- Ehlert, F.J., Griffin, M.T., Sawyer, G.W., and Bailon, R. (1999). A simple method for estimation of agonist activity at receptor subtypes: comparison of native and cloned M3 muscarinic receptors in guinea pig ileum and transfected cells. *J. Pharmacol. Exp. Ther.* **289**, 981–992.
- Finn, R.D., Coghill, P., Eberhardt, R.Y., Eddy, S.R., Mistry, J., Mitchell, A.L., Potter, S.C., Punta, M., Qureshi, M., Sangrador-Vegas, A., et al. (2016). The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res.* **44** (D1), D279–D285.
- Flock, T., Ravarani, C.N.J., Sun, D., Venkatakrisnan, A.J., Kayikci, M., Tate, C.G., Veprintsev, D.B., and Babu, M.M. (2015). Universal allosteric mechanism for G α activation by GPCRs. *Nature* **524**, 173–179.
- Flock, T., Hauser, A.S., Lund, N., Gloriam, D.E., Balaji, S., and Babu, M.M. (2017). Selectivity determinants of GPCR-G-protein binding. *Nature* **545**, 317–322.
- Galés, C., Rebois, R.V., Hogue, M., Trieu, P., Breit, A., Hébert, T.E., and Bouvier, M. (2005). Real-time monitoring of receptor and G-protein interactions in living cells. *Nat. Methods* **2**, 177–184.
- García-Nafria, J., Lee, Y., Bai, X., Carpenter, B., and Tate, C.G. (2018a). Cryo-EM structure of the adenosine A_{2A} receptor coupled to an engineered heterotrimeric G protein. *eLife* **7**, 7.
- García-Nafria, J., Nehmé, R., Edwards, P.C., and Tate, C.G. (2018b). Cryo-EM structure of the serotonin 5-HT_{1B} receptor coupled to heterotrimeric G_o. *Nature* **558**, 620–623.
- Guettier, J.M., Gautam, D., Scarselli, M., Ruiz de Azua, I., Li, J.H., Rosemond, E., Ma, X., Gonzalez, F.J., Armbruster, B.N., Lu, H., et al. (2009). A chemical-genetic approach to study G protein regulation of beta cell function in vivo. *Proc. Natl. Acad. Sci. USA* **106**, 19197–19202.
- Hannenhalli, S.S., and Russell, R.B. (2000). Analysis and prediction of functional sub-types from protein sequence alignments. *J. Mol. Biol.* **303**, 61–76.
- Harding, S.D., Sharman, J.L., Faccenda, E., Southan, C., Pawson, A.J., Ireland, S., Gray, A.J.G., Bruce, L., Alexander, S.P.H., Anderton, S., et al.; NC-IUPHAR (2018). The IUPHAR/BPS Guide to PHARMACOLOGY in 2018: updates and expansion to encompass the new guide to IMMUNOPHARMACOLOGY. *Nucleic Acids Res.* **46** (D1), D1091–D1106.
- Hauser, A.S., Attwood, M.M., Rask-Andersen, M., Schiöth, H.B., and Gloriam, D.E. (2017). Trends in GPCR drug discovery: new agents, targets and indications. *Nat. Rev. Drug Discov.* **16**, 829–842.
- Hauser, A.S., Chavali, S., Masuho, I., Jahn, L.J., Martemyanov, K.A., Gloriam, D.E., and Babu, M.M. (2018). Pharmacogenomics of GPCR Drug Targets. *Cell* **172**, 41–54.
- Herroeder, S., Reichardt, P., Sassmann, A., Zimmermann, B., Jaeneke, D., Hoeckner, J., Hollmann, M.W., Fischer, K.D., Vogt, S., Grosse, R., et al. (2009). Guanine nucleotide-binding proteins of the G12 family shape immune functions by controlling CD4+ T cell adhesiveness and motility. *Immunity* **30**, 708–720.
- Horn, F., van der Wenden, E.M., Oliveira, L., IJzerman, A.P., and Vriend, G. (2000). Receptors coupling to G proteins: is there a signal behind the sequence? *Proteins* **41**, 448–459.
- Inoue, A., Ishiguro, J., Kitamura, H., Arima, N., Okutani, M., Shuto, A., Higashiyama, S., Ohwada, T., Arai, H., Makide, K., and Aoki, J. (2012). TGF α shedding assay: an accurate and versatile method for detecting GPCR activation. *Nat. Methods* **9**, 1021–1029.
- Insel, P.A., and Ostrom, R.S. (2003). Forskolin as a tool for examining adenylyl cyclase expression, regulation, and G protein signaling. *Cell. Mol. Neurobiol.* **23**, 305–314.
- Isberg, V., Mordalski, S., Munk, C., Rataj, K., Harpsøe, K., Hauser, A.S., Vroiling, B., Bojarski, A.J., Vriend, G., and Gloriam, D.E. (2017). GPCRdb: an information system for G protein-coupled receptors. *Nucleic Acids Res.* **45**, 2936.
- Kang, Y., Kuybeda, O., de Waal, P.W., Mukherjee, S., Van Eps, N., Dutka, P., Zhou, X.E., Bartesaghi, A., Erramilli, S., Morizumi, T., et al. (2018). Cryo-EM structure of human rhodopsin bound to an inhibitory G protein. *Nature* **558**, 553–558.
- Kihara, Y., Maceyka, M., Spiegel, S., and Chun, J. (2014). Lysophospholipid receptor nomenclature review: IUPHAR Review 8. *Br. J. Pharmacol.* **171**, 3575–3594.
- Koehl, A., Hu, H., Maeda, S., Zhang, Y., Qu, Q., Paggi, J.M., Latorraca, N.R., Hilger, D., Dawson, R., Matile, H., et al. (2018). Structure of the μ -opioid receptor-G_i protein complex. *Nature* **558**, 547–552.
- Leng, W., Pang, X., Xia, H., Li, M., Chen, L., Tang, Q., Yuan, D., Li, R., Li, L., Gao, F., and Bi, F. (2013). Novel split-luciferase-based genetically encoded biosensors for noninvasive visualization of Rho GTPases. *PLoS ONE* **8**, e62230.
- Martin, D., Abba, M.C., Molinolo, A.A., Vitale-Cross, L., Wang, Z., Zaida, M., Delic, N.C., Samuels, Y., Lyons, J.G., and Gutkind, J.S. (2014). The head and neck cancer cell oncogene: a platform for the development of precision molecular therapies. *Oncotarget* **5**, 8906–8923.
- Muppidi, J.R., Schmitz, R., Green, J.A., Xiao, W., Larsen, A.B., Braun, S.E., An, J., Xu, Y., Rosenwald, A., Ott, G., et al. (2014). Loss of signalling via G α 13 in germinal centre B-cell-derived lymphoma. *Nature* **516**, 254–258.
- O'Hayre, M., Inoue, A., Kufareva, I., Wang, Z., Mikelis, C.M., Drummond, R.A., Avino, S., Finkel, K., Kalim, K.W., DiPasquale, G., et al. (2016). Inactivating mutations in GNA13 and RHOA in Burkitt's lymphoma and diffuse large B-cell

- lymphoma: a tumor suppressor function for the $G\alpha_{13}/RhoA$ axis in B cells. *Oncogene* 35, 3771–3780.
- Papaleo, E., Saladino, G., Lambrugh, M., Lindorff-Larsen, K., Gervasio, F.L., and Nussinov, R. (2016). The Role of Protein Loops and Linkers in Conformational Dynamics and Allostery. *Chem. Rev.* 116, 6391–6423.
- Patel, Y.M., Lordkipanidzé, M., Lowe, G.C., Nisar, S.P., Garner, K., Stockley, J., Daly, M.E., Mitchell, M., Watson, S.P., Austin, S.K., and Mundell, S.J. (2014). A novel mutation in the P2Y₁₂ receptor and a function-reducing polymorphism in protease-activated receptor 1 in a patient with chronic bleeding. *J. Thromb. Haemost.* 12, 716–725.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011). Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* 12, 2825–2830.
- Raimondi, F., Felling, A., Portella, G., Orozco, M., and Fanelli, F. (2013a). Light on the structural communication in Ras GTPases. *J. Biomol. Struct. Dyn.* 31, 142–157.
- Raimondi, F., Felling, A., Seeber, M., Mariani, S., and Fanelli, F. (2013b). A Mixed Protein Structure Network and Elastic Network Model Approach to Predict the Structural Communication in Biomolecular Systems: The PDZ2 Domain from Tyrosine Phosphatase 1E As a Case Study. *J. Chem. Theory Comput.* 9, 2504–2518.
- Rasmussen, S.G., DeVree, B.T., Zou, Y., Kruse, A.C., Chung, K.Y., Kobilka, T.S., Thian, F.S., Chae, P.S., Pardon, E., Calinski, D., et al. (2011). Crystal structure of the β_2 adrenergic receptor-Gs protein complex. *Nature* 477, 549–555.
- Rodriguez, G.J., Yao, R., Lichtarge, O., and Wensel, T.G. (2010). Evolution-guided discovery and recoding of allosteric pathway specificity determinants in psychoactive bioamine receptors. *Proc. Natl. Acad. Sci. USA* 107, 7787–7792.
- Salzberg, S.L. (1997). On Comparing Classifiers: Pitfalls to Avoid and a Recommended Approach. *Data Min. Knowl. Discov.* 1, 317–328.
- Saulière, A., Bellot, M., Paris, H., Denis, C., Finana, F., Hansen, J.T., Alté, M.F., Seguelas, M.H., Pathak, A., Hansen, J.L., et al. (2012). Deciphering biased-agonism complexity reveals a new active AT₁ receptor entity. *Nat. Chem. Biol.* 8, 622–630.
- Schrage, R., Schmitz, A.L., Gaffal, E., Annala, S., Kehraus, S., Wenzel, D., Bülesbach, K.M., Bald, T., Inoue, A., Shinjo, Y., et al. (2015). The experimental power of FR900359 to study Gq-regulated biological processes. *Nat. Commun.* 6, 10156.
- Seeber, M., Felling, A., Raimondi, F., Mariani, S., and Fanelli, F. (2015). WebPSN: a web server for high-throughput investigation of structural communication in biomacromolecules. *Bioinformatics* 31, 779–781.
- Sgourakis, N.G., Bagos, P.G., and Hamodrakas, S.J. (2005a). Prediction of the coupling specificity of GPCRs to four families of G-proteins using hidden Markov models and artificial neural networks. *Bioinformatics* 21, 4101–4106.
- Sgourakis, N.G., Bagos, P.G., Papsaikas, P.K., and Hamodrakas, S.J. (2005b). A method for the prediction of GPCRs coupling specificity to G-proteins using refined profile Hidden Markov Models. *BMC Bioinformatics* 6, 104.
- Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., Söding, J., et al. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* 7, 539.
- Stallaert, W., van der Westhuizen, E.T., Schönegge, A.M., Plouffe, B., Hogue, M., Lukashova, V., Inoue, A., Ishida, S., Aoki, J., Le Guillou, C., and Bouvier, M. (2017). Purinergic Receptor Transactivation by the β_2 -Adrenergic Receptor Increases Intracellular Ca^{2+} in Nonexcitable Cells. *Mol. Pharmacol.* 91, 533–544.
- Sugimoto, Y., and Narumiya, S. (2007). Prostaglandin E receptors. *J. Biol. Chem.* 282, 11613–11617.
- Suzuki, N., Hajicek, N., and Kozasa, T. (2009). Regulation and physiological functions of G12/13-mediated signaling pathways. *Neurosignals* 17, 55–70.
- Thomsen, W., Frazer, J., and Unett, D. (2005). Functional assays for screening GPCR targets. *Curr. Opin. Biotechnol.* 16, 655–665.
- Urban, D.J., and Roth, B.L. (2015). DREADDs (designer receptors exclusively activated by designer drugs): chemogenetic tools with therapeutic utility. *Annu. Rev. Pharmacol. Toxicol.* 55, 399–417.
- Van Eps, N., Altenbach, C., Caro, L.N., Latorraca, N.R., Hollingsworth, S.A., Dror, R.O., Ernst, O.P., and Hubbell, W.L. (2018). G_i- and G_s-coupled GPCRs show different modes of G-protein binding. *Proc. Natl. Acad. Sci. USA* 115, 2383–2388.
- Velankar, S., Dana, J.M., Jacobsen, J., van Ginkel, G., Gane, P.J., Luo, J., Oldfield, T.J., O'Donovan, C., Martin, M.J., and Kleywegt, G.J. (2013). SIFTS: Structure Integration with Function, Taxonomy and Sequences resource. *Nucleic Acids Res.* 41, D483–D489.
- Venkatakrishnan, A.J., Deupi, X., Lebon, G., Tate, C.G., Schertler, G.F., and Babu, M.M. (2013). Molecular signatures of G-protein-coupled receptors. *Nature* 494, 185–194.
- Venkatakrishnan, A.J., Flock, T., Prado, D.E., Oates, M.E., Gough, J., and Madan Babu, M. (2014). Structured and disordered facets of the GPCR fold. *Curr. Opin. Struct. Biol.* 27, 129–137.
- Venkatakrishnan, A.J., Deupi, X., Lebon, G., Heydenreich, F.M., Flock, T., Miljus, T., Balaji, S., Bouvier, M., Veprincev, D.B., Tate, C.G., et al. (2016). Diverse activation pathways in class A GPCRs converge near the G-protein-coupling region. *Nature* 536, 484–487.
- Violin, J.D., Crombie, A.L., Soergel, D.G., and Lark, M.W. (2014). Biased ligands at G-protein-coupled receptors: promise and progress. *Trends Pharmacol. Sci.* 35, 308–316.
- Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., and Barton, G.J. (2009). Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25, 1189–1191.
- Weinstein, J.B.H. (1995). Integrated methods for the construction of three-dimensional models and computational probing of structure-function relations in G protein-coupled receptors. *Methods Neurosci.* 25, 366–428.
- Wess, J., Nakajima, K., and Jain, S. (2013). Novel designer receptors to probe GPCR signaling and physiology. *Trends Pharmacol. Sci.* 34, 385–392.
- Wettschureck, N., and Offermanns, S. (2005). Mammalian G proteins and their cell type specific functions. *Physiol. Rev.* 85, 1159–1204.
- Wheeler, T.J., Clements, J., and Finn, R.D. (2014). Skylign: a tool for creating informative, interactive logos representing sequence alignments and profile hidden Markov models. *BMC Bioinformatics* 15, 7.
- Wong, S.K. (2003). G protein selectivity is regulated by multiple intracellular regions of GPCRs. *Neurosignals* 12, 1–12.
- Woodward, D.F., Jones, R.L., and Narumiya, S. (2011). International Union of Basic and Clinical Pharmacology. LXXXIII: classification of prostanoid receptors, updating 15 years of progress. *Pharmacol. Rev.* 63, 471–538.
- Yabuki, Y., Muramatsu, T., Hirokawa, T., Mukai, H., and Suwa, M. (2005). GRIFFIN: a system for predicting GPCR-G-protein coupling selectivity using a support vector machine and a hidden Markov model. *Nucleic Acids Res.* 33, W148–W153.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti DYKDDDDK tag antibody (mouse monoclonal, clone 1E6)	FujiFilm Wako Pure Chemicals	Cat# 012-22384; RRID: AB_10659717
Goat anti-mouse IgG secondary antibody conjugated with Alexa Fluor 488	Thermo Fisher Scientific	Cat# A28175; RRID: AB_2536161
Anti-G α q antibody (goat polyclonal)	Abcam	Cat# ab128060; RRID: AB_11145302
Anti-G α 11 antibody (mouse monoclonal, clone D-6; recognizing G α q and G α 11)	Santa Cruz Biotechnologies	Cat# sc-390382; RRID: AB_2715558
Anti-G α 13 antibody (rabbit monoclonal, clone EPR5436)	Abcam	Cat# ab128900; RRID: AB_11142234
Anti- α -tubulin antibody (mouse monoclonal, clone DM1A)	Santa Cruz Biotechnologies	Cat# sc-32293; RRID: AB_628412
Anti-goat IgG secondary antibody conjugated with horseradish peroxidase (HRP)	American Qualex	A201PS
Anti-mouse IgG secondary antibody conjugated with horseradish peroxidase (HRP)	GE Healthcare	Cat# NA9310; RRID: AB_772193
Anti-rabbit IgG secondary antibody conjugated with horseradish peroxidase (HRP)	GE Healthcare	Cat# NA9340; RRID: AB_772191
Anti-RhoA antibody (mouse monoclonal)	Cytoskeleton	Cat# ARH04; RRID: AB_2728698
Bacterial and Virus Strains		
SCS1 Supercompetent Cells	Agilent Technologies	200231
Biological Samples		
Fetal bovine serum	Thermo Fisher Scientific	10270-106
Fetal bovine serum (for HN12 cells and Cal27 cells)	Sigma-Aldrich	F2442
Goat serum	Nippon Bio-test Laboratories	0208-01
Skim milk powder	FUJIFILM Wako Pure Chemical	198-10605
Chemicals, Peptides, and Recombinant Proteins		
Lipofectamine 2000	Thermo Fisher Scientific	11668-019
Lipofectamine RNAiMax	Thermo Fisher Scientific	13778-150
Polyethylenimine (PEI) Max, (Mw 40,000)	Polysciences	24765-1
Opti-MEM	Thermo Fisher Scientific	31985-070
HEPES	Sigma-Aldrich	H3375
EDTA solution	Thermo Fisher Scientific	15575-020
<i>p</i> -nitrophenylphosphate, disodium salt	FUJIFILM Wako Pure Chemical	145-02344
Trypsin	Thermo Fisher Scientific	27250-018
Bovine serum albumin	SERVA Electrophoresis	11945
Coelenterazine	Carbosynth	EC14031
D-luciferin, potassium salt	FUJIFILM Wako Pure Chemical	126-05116
2-oxo-glutaric acid	FUJIFILM Wako Pure Chemical	117-00081
7 α ,25-dihydroxycholesterol	Cayman Chemical	11032
9-HODE	Cayman Chemical	38400
Acetylcholine	Sigma-Aldrich	A6625
Adenosine	FUJIFILM Wako Pure Chemical	011-24593
ADP	FUJIFILM Wako Pure Chemical	015-13241
Angiotensin II	Peptide Institute	4001-v
Apelin	Cayman Chemical	13523
ATP	FUJIFILM Wako Pure Chemical	017-21511
BAM-22P	Anygen	AGP-8605

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bombesin	Peptide Institute	4086-v
Bradykinin	Peptide Institute	4002-v
Capric acid	FUJIFILM Wako Pure Chemical	041-23256
CAY10583	Cayman Chemical	10012424
CCK-Octapeptide	Peptide Institute	4100-v
Cortistatin-14	Peptide Institute	4329-v
CP-55940	Cayman Chemical	90084
Dopamine	FUJIFILM Wako Pure Chemical	040-15433
Dynorphin A	Peptide Institute	4080-v
Endothelin-1	Peptide Institute	4198 s
fMLP	Peptide Institute	4066-v
Galanin	Abgent	SP2781a
Galanin (rat)	Peptide Institute	4244-v
Gastric Inhibitory Peptide	Abgent	SP3260a
GSK1292263	AdooQ BioScience	A10439-10
Histamine	FUJIFILM Wako Pure Chemical	087-03553
Iloprost	Cayman Chemical	18215
Isoproterenol	Sigma-Aldrich	I6504
Kisspeptin-10	Peptide Institute	4389-v
Leukotriene B4	Cayman Chemical	20110
Leukotriene D4	Cayman Chemical	20310
Leuprolide	GenScript	RP17321
Lysophosphatidic acid	Avanti Polar Lipids	857130
Lysophosphatidylinositol	Avanti Polar Lipids	850105
Lysophosphatidylserine	Avanti Polar Lipids	858143
Lysyl-Bradykinin	Peptide Institute	4008-v
MCH	Peptide Institute	4369-v
MDL 29951	Cayman Chemical	16266
Melatonin	FUJIFILM Wako Pure Chemical	135-17113
Methionine-Enkephalin	Peptide Institute	4042-v
Methionine-Enkephalin	Peptide Institute	4042-v
MK-0677	Cayman Chemical	18003
Motilin	GenScript	RP10527
Neurokinin A	Peptide Institute	4154-v
Neurokinin B	Peptide Institute	4317-v
Neuromedin B	Peptide Institute	4152-v
Neuromedin-U25	Peptide Institute	4490-v
Neuropeptide FF	GenScript	RP10470
Neuropeptide W-30	Abgent	SP2086a
Nociceptin	Peptide Institute	4318-v
Norepinephrine	Sigma-Aldrich	N5785
Orexin-A	Peptide Institute	4346 s
Oxytocin	Peptide Institute	4084-v
PACAP27	Peptide Institute	4231-v
PAF	Avanti Polar Lipids	878110
Palmitic acid	FUJIFILM Wako Pure Chemical	165-00102
PAR1 peptide	Peptide Institute	PAR-3665-PI
PAR2 peptide	Peptide Institute	PAR-3664-PI

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Parathyroid Hormone	Peptide Institute	4068 s
Prolactin-Releasing Peptide-20	Anygen	AGP-8642
Propionic acid	FUJIFILM Wako Pure Chemical	163-04726
Prostaglandin D2	Cayman Chemical	12010
Prostaglandin E2	Cayman Chemical	14010
Prostaglandin F2 α	Cayman Chemical	16010
Serotonin	FUJIFILM Wako Pure Chemical	321-42341
Somatostatin	Peptide Institute	4023-v
Sphingosine 1-phosphate	Cayman Chemical	62570
Substance P	Peptide Institute	4014-v
Thrombin	Sigma-Aldrich	T4648
TIP39	Peptide Institute	4479 s
TUG 891	Tocris Bioscience	4601
U-46619	Cayman Chemical	16450
UDP	FUJIFILM Wako Pure Chemical	212-00861
UDP-Glucose	FUJIFILM Wako Pure Chemical	214-00941
Urotensin II	Abgent	SP2662a
Vasopressin	Peptide Institute	4085-v
WKYMVM	Anygen	AGP-8694
Zaprinast	Cayman Chemical	10010421
α -MSH	Peptide Institute	4057-v
SII	Peptide Institute	4476-v
Clozapine N-oxide	Tocris Bioscience	4936
KI16425	AdooQ BioScience	A10501
JTE-013	Cayman Chemical	10009458
Forskolin	FUJIFILM Wako Pure Chemical	067-02191
DMEM	Nissui	05919
Penicillin G	Sigma-Aldrich	P3032
Streptomycin	Thermo Fisher Scientific	11860-038
Probenecid	FUJIFILM Wako Pure Chemical	162-26112
ImmunoStar Zeta	FUJIFILM Wako Pure Chemical	291-72401
Angiotensin II (for RhoA pulldown assay)	Sigma-Aldrich	A9525
CP-55940 (for RhoA pulldown assay)	Sigma-Aldrich	C1112
Lysophosphatidic acid (for RhoA pulldown assay)	Sigma-Aldrich	L7260
Luminol Sodium Salt HG	FUJIFILM Wako Pure Chemical	123-04021
Critical Commercial Assays		
PrimeSTAR HS DNA Polymerase	Takara Bio	R010B
NEBuilder HiFi DNA Assembly Master Mix	New England BioLabs	E2621X
NucleoBond Xtra Midi Plus	Macherey-Nagel	740412.50
PureYield Plasmid Miniprep System	Promega	A1222
MycAlert Mycoplasma Detection Kit	Lonza	LT07-118
High-Capacity cDNA Reverse Transcription Kits	Thermo Fisher Scientific	4368814
SYBR Premix Ex Taq	Takara Bio	RR041A
FLIPR Calcium 6 Assay Kit	Molecular Devices	N/A
RhoA Pull-Down Activation Assay Biochem Kit	Cytoskeleton	BK036
DC Protein Assay	BioRad	5000112
Experimental Models: Cell Lines		
Human: HEK293 cells (parental HEK293)	Inoue et al., 2012	N/A
Δ Gq HEK293 cells	Schrage et al., 2015	CL1

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Δ G12 HEK293 cells	Devost et al., 2017	CL3
Δ Gq/ Δ G12 HEK293 cells	Devost et al., 2017	CL4-1
Δ Gs HEK293 cells	Stallaert et al., 2017	CL4
MDA-MB-231 cells	ATCC	HTB26
PC-3 cells	ATCC	CRL1435
HN12 cells	NIH cell collection	N/A
Cal27 cells	NIH cell collection	N/A
Oligonucleotides		
siRNA targeting sequence: GNAQ (#1) 5'-GGAGAGAGU GGCAAGAGUACGUUUU-3'	Thermo Fisher Scientific	GNAQHSS104236
siRNA targeting sequence: GNAQ (#2) 5'-CCCUUUGAC UUACAAAGUGUCAUUU-3'	Thermo Fisher Scientific	GNAQHSS104237
siRNA targeting sequence: GNA11 (#1) 5'-CCGGCAUC AUCGAGUACCCUUUCGA-3'	Thermo Fisher Scientific	GNA11HSS178464
siRNA targeting sequence: GNA11 (#2) 5'-GCAUCAGU ACGUCAGUGCCAUCAAG-3'	Thermo Fisher Scientific	GNA11HSS104213
siRNA targeting sequence: GNA12 (#1) 5'-CCAAGGGA AUUGUGGAGCAUGACUU-3'	Thermo Fisher Scientific	GNA12-HSS178466
siRNA targeting sequence: GNA12 (#2) 5'-CCAUCGUC AACACAAGCUCUUCUU-3'	Thermo Fisher Scientific	GNA12MSS204749
siRNA targeting sequence: GNA13 (#1) 5'-CAGAAGCC CUUAUACCACCACUUCA-3'	Thermo Fisher Scientific	GNA13-HSS173827
siRNA targeting sequence: GNA13 (#2) 5'-GCAGCCCAA GGAAUGGUGGAAACAA-3'	Thermo Fisher Scientific	GNA13-HSS116479
siRNA targeting sequence: ADAM17 5'-CAGAAUCGU GUUGACAGCAAAGAAA-3'	Thermo Fisher Scientific	ADAM17-HSS186181
Stealth siRNA control	Thermo Fisher Scientific	12935113
qPCR Primers for the GNAQ gene: 5'-ACCGAAT GGAGGAAAGCAAGG-3' 5'-CATCTCTCTGGGGTCC ATCATATTC-3'	This paper	N/A
qPCR Primers for the GNA11 gene: 5'-CAGCGAATACG ACCAAGTCC-3' 5'-ACCAGGGGTAGGTGATGATG-3'	This paper	N/A
qPCR Primers for the GNA12 gene: 5'-GAGGGATTCT GGCATCAGG-3' 5'-CGATCCGGTCCAAGTTGTC-3'	This paper	N/A
qPCR Primers for the GNA13 gene: 5'-CCTGGATAACT TGGATAAACTTGG-3' 5'-TTCATGGATGCCTTTGGTG-3'	This paper	N/A
qPCR Primers for the GAPDH gene: 5'-GCCAAGGTCA TCCATGACAACT-3' 5'-GAGGGGCCATCCACAGTCTT-3'	This paper	N/A
Recombinant DNA		
Human GPCR-encoding plasmids library	Inoue et al., 2012 and this study	N/A
AP-TGF- α (codon-optimized)	Inoue et al., 2012	N/A
Chimeric G α subunits	Inoue et al., 2012 and this study	N/A
NanoBiT-G-proteins	This study	N/A
NanoBiT-RhoA sensor	This study	N/A
NanoBiT-IP3 sensor	This study	N/A
M3D	This study	N/A
M4D	This study	N/A
M3D-Gs (codon-optimized)	This study	N/A
ICL3-swapped M3D constructs	This study	N/A
ICL3- and C-terminus-swapped M3D constructs	This study	N/A
M3D-GPR183/ICL3	This study	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
M3D-GPR132/ICL3	This study	N/A
Glo-22F (codon-optimized)	This study	N/A
Software and Algorithms		
PyMol, Version 1.7.2	N/A	https://pymol.org
Cytoscape, Version 3.5	N/A	https://cytoscape.org/
GraphPad Prism, Version 7	GraphPad	N/A
Soft Max Pro, Version 7	Molecular Devices	N/A
FlowJo software, Version 10	FlowJo	N/A
Multi Gauge, Version 3.0	Fujifilm	N/A
HMMER3, Version 3.1b2	Eddy, Bioinformatics, 1998	http://www.hmmer.org/
Numpy, Version 1.11	N/A	http://www.numpy.org/
Scipy, Version 0.18	N/A	https://www.scipy.org
scikit-learn, Version 0.18	Pedregosa et al., 2011	https://www.scikit-learn.org
igraph, Version 0.7	Csardi and Nepusz, 2006	https://igraph.org/python/
Other		
96W clear plate for cells	Greiner Bio-one	655180
96W clear plate for conditioned media	Greiner Bio-one	655101
96W white plate	Greiner Bio-one	655083
96W white plate, half-area	Greiner Bio-one	675083
96W black plate, half-area, clear-bottom	Greiner Bio-one	675986
Absorbance microplate reader	Molecular Devices	SpectraMax 340PC384
Microplate stacker	Molecular Devices	StakMax
Luminescence microplate reader	Molecular Devices	SpectraMax L
Multi-mode, auto-pipetting microplate reader	Molecular Devices	FlexStation 3
Flow cytometer	Sony	EC800
Chemiluminescence imager	Fujifilm	LAS-4000
An online resource for the G-protein-coupling data and the prediction tool	This study	gpcr.russelllab.org

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled upon reasonable request by the Lead Contact, Asuka Inoue (iaska@tohoku.ac.jp).

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Cells and transfection**

HEK293A cells (Female origin; Thermo Fisher Scientific) and their derivative G-protein-deficient HEK293 cells were maintained in Dulbecco's Modified Eagle Medium (DMEM 2, Nissui Pharmaceutical) supplemented with 10% fetal bovine serum (GIBCO, Thermo Fisher Scientific) and penicillin-streptomycin-glutamine (complete DMEM). Generation and characterization of the ΔG_q HEK293 cells, in which null mutations were introduced into the *GNAQ* and the *GNA11* genes by a CRISPR-Cas9 system ([Schrage et al., 2015](#)) and thus their functional products are lacking, the ΔG_{12} HEK293 cells (lacking functional products of the *GNA12* and the *GNA13* genes), the $\Delta G_q/\Delta G_{12}$ HEK293 cells (lacking those of the *GNAQ*, the *GNA11*, *GNA12* and the *GNA13* genes) ([Devost et al., 2017](#)) and the ΔG_s HEK293 cells (lacking those of the *GNAS* and the *GNAL* genes) ([Stallaert et al., 2017](#)) were described previously. The cells were regularly tested for mycoplasma contamination using a MycoAlert Mycoplasma Detection Kit (Lonza).

Transfection was performed by using a lipofection reagent, Lipofectamine® 2000 Reagent (Thermo Fisher Scientific), or polyethylenimine (PEI) solution (Polyethylenimine "Max," Polysciences). Typically, HEK293 cells were seeded in a 6-well culture plate at cell density of 2×10^5 cells ml^{-1} in 2 mL of the complete DMEM and cultured for one day in a humidified 37°C incubator with 5% CO₂. Seeding density for the ΔG_{12} cells and the $\Delta G_q/\Delta G_{12}$ cells were increased to 2.5×10^5 cells ml^{-1} owing to slower growth of the cells

than the parent HEK293 cells and the ΔG_q cells. For Lipofectamine® 2000 transfection, a transfection mixture was prepared by mixing plasmid solution diluted in 250 μ l of Opti-MEM (Life Technologies) and Lipofectamine® 2000 solution (2.5 μ l) in 250 μ l of Opti-MEM. For PEI transfection, a transfection solution was mixed by combining plasmid solution diluted in 100 μ l of Opti-MEM and 4 μ l of 1 mg ml⁻¹ PEI solution in 100 μ l of Opti-MEM. Both Lipofectamine® 2000 and the PEI transfection gave almost identical transfection efficiency in our culture condition. The transfected cells were further incubated for one day before subjected to an assay as described below.

MDA-MB-231 cells (female origin) and PC-3 cells (male origin) were maintained in RPMI 1640 (Nissui Pharmaceutical) supplemented with 5% fetal bovine serum and penicillin-streptomycin-glutamine. MDA-MB-231 cells and PC-3 cells were seeded in a 10-cm culture dish at cell density of 2×10^5 cells ml⁻¹ in 10 mL of the media and cultured for one day in the incubator. Transfection was performed by using 20 μ l of Lipofectamine® 2000 transfection reagent. The transfected cells were incubated for one day before subjected to the NanoBiT-RhoA assay as described below.

HN12 cells (female origin) and Cal27 cells (male origin), which were characterized as part of a head and neck cancer cell oncogene effort (Martin et al., 2014) and obtained from this NIH cell collection, were maintained in DMEM supplemented with 10% FBS (Sigma-Aldrich).

METHOD DETAILS

Plasmids

Only human GPCRs and human $G\alpha$ subunits were used in this study. An open reading frame of each full-length GPCR was cloned into pCAGGS expression plasmid (a kind gift from Dr. Jun-ichi Miyazaki at Osaka University, Japan) or pcDNA3.1 expression plasmid. Except when otherwise specified, GPCR sequences were devoid of epitope tags. The GPCRs examined for this study (148 GPCRs) originated from a previous GPCR library (109 GPCRs) (Inoue et al., 2012) and an extended list of GPCR families (39 GPCRs). In our library, we covered all of the members for selected GPCR families. We note that there are 8 GPCRs (AGTR2, GPBAR1, GPER, GPR18, HTR5A, MC2R, NPBWR2 and PTGDR2) that were unresponsive in the chimeric G-protein-based TGF- α shedding assay (data not shown) and thus were not included in the G-protein-coupling dataset.

Full-length, untagged $G\alpha$ subunits were cloned into the pCAGGS plasmid. Chimeric $G\alpha$ subunits, in which the C-terminal 6 amino acids were substituted, were generated with PCR-amplified fragments using synthesized oligonucleotides encoding swapped C-terminal sequences. A C-terminally truncated $G\alpha_q$ subunit, which lacked the 7 amino acids (note that the -7 position is identical among all of the $G\alpha$ subunits), was used as a negative control for the chimeric-G-protein-based TGF- α shedding assay. Inserted sequences were verified by Sanger sequencing (Fasmac). Codon-optimized AP-TGF- α cloned into the pCAGGS plasmid was used in this study. Amino acid sequences for the AP-TGF- α construct and the chimeric $G\alpha$ subunits are listed in [Data S3](#).

M3D and M4D (Armbruster et al., 2007) were generated by introducing the two mutations ($Y^{3.33}C$ and $A^{5.46}G$), which alter ligand specificity from ACh to CNO, in human CHRM3 (corresponding to Y149C and A239G) and CHRM4 (Y113C and A203G), respectively, by using an NEBuilder HiFi DNA Assembly system (New England Biolabs) and cloned into the pcDNA3.1 vector with N-terminal FLAG-epitope (DYKDDDDK) tag. ICL3-substituted M3D chimeras were constructed by the NEBuilder system with PCR-amplified fragments using synthesized oligonucleotides encoding swapped ICL3 sequences. Dual ICL3- and C-terminally-substituted M3D chimeras were generated by assembling PCR-amplified fragments with synthesized oligonucleotides for C-terminal sequences. The substituted ICL3 and C-terminus correspond to nucleotide positions of 778-1455 and 1633-1770 of the CHRM3 ORF. A coding sequence for M3D- G_s (Guettier et al., 2009) was human codon-optimized and gene-synthesized by Genscript and inserted into pcDNA3.1 with the N-terminal FLAG-epitope tag. Throughout the study, we used the same N-terminally FLAG-tagged DREADD constructs for functional assays and expression analysis. DNA sequences and translated amino acid sequences for the DREADDs are shown in [Data S3](#).

For NanoBiT-G-proteins, the large fragment (LgBiT) of the NanoBiT luciferase was inserted into the helical domain of human $G\alpha$ subunit ($G\alpha$ -Lg) flanked by 15-amino acid flexible linkers (GGSGGGSGGSSSGG) and the small fragment (SmBiT) was N-terminally fused to human $G\beta$ subunit (Sm- $G\beta$) or human $G\gamma$ subunit (Sm- $G\gamma$) with the 15-amino acid linker. A coding sequence for the $G\alpha$ -Lg was human codon-optimized and gene-synthesized by Genscript and inserted into pcDNA3.1 plasmid. To construct a coding sequence for the Sm- $G\beta$ and the Sm- $G\gamma$, oligonucleotides encoding the N-terminal SmBiT-linker (Fasmac) and PCR-amplified fragment of full-length $G\beta$ ($G\beta_1$, $G\beta_3$ or $G\beta_5$) or $G\gamma$ ($G\gamma_2$ or $G\gamma_{11}$) were assembled by using the NEBuilder system and cloned into the pCAGGS vector. Coding sequences for untagged $G\beta_1$ and $G\gamma_2$ were inserted into pcDNA3.1 vector. Coding sequences for *RIC8A* and *RIC8B* (isoform 2) were cloned into pCAGGS vector. Amino acid sequences for the NanoBiT-G-proteins are listed in [Data S3](#).

For generation of the NanoBiT-RhoA sensor, we replaced firefly luciferase fragments of previously described RhoA constructs (Leng et al., 2013) with the NanoBiT fragments. Specifically, LgBiT and SmBiT were N-terminally fused to human RhoA (residues 2-193) and the GTPase-binding domain (GBD) of human PKN1 (residues 13-112), a RhoA effector, respectively, with the 15-amino acid linker. A coding sequence for RhoA and PKN1-GBD was human codon-optimized and gene-synthesized by Genscript and inserted into the pCAGGS plasmid by following a similar method as described in the NanoBiT-G-protein construction. Amino acid sequences for the NanoBiT-RhoA constructs (Lg-RhoA and Sm-PKN1) are listed in [Data S3](#).

Similarly, to construct the NanoBiT-IP₃ sensor, we exchanged firefly luciferase fragments of a previously described IP₃ construct (Ataei et al., 2013) with the NanoBiT fragments. Specifically, LgBiT and SmBiT were fused to N-terminus and C-terminus, respectively, of IP₃-binding core domain (IBC) of human type 2 IP₃ receptor (Gene symbol ITPR2; residues 225-604), flanked by the 15-amino acid linker. A coding sequence for ITPR2-IBC was human codon-optimized and gene-synthesized by Genscript, and inserted into the pCAGGS plasmid by following the above-described method. An amino acid sequence for the NanoBiT-IP₃ sensor (Lg-IP3R2-Sm) is listed in [Data S3](#).

TGF- α shedding assay

The TGF- α shedding assay was performed as described previously (Inoue et al., 2012) with minor modifications. Plasmid transfection was performed in a 6-well plate with a mixture of 500 ng AP-TGF- α -encoding plasmid, 200 ng GPCR-encoding plasmid with or without 100 ng G α -encoding plasmid (per well, hereafter). After 1-day culture, the transfected cells were harvested by trypsinization, pelleted by centrifugation at 190 g for 5 min and washed once with Hank's Balanced Salt Solution (HBSS) containing 5 mM HEPES (pH 7.4). After centrifugation, the cells were resuspended in 6 mL of the HEPES-containing HBSS. We note that trypsinization and following washing procedure resulted in a higher signal-to-background TGF- α shedding response as compared with harvesting cells without trypsin (EDTA only). The cell suspension was seeded in a 96-well culture plate (cell plate) at a volume of 90 μ l (per well hereafter) and incubated for 30 min in a 5% CO₂ incubator at 37°C. The cells were treated with a GPCR ligand (10X, diluted in HBSS containing 5 mM HEPES (pH 7.4) and 0.01% (w/v) bovine serum albumin (BSA, fatty acid-free and protease-free grade; Serva)). After spinning the cell plates, conditioned media (80 μ l) was transferred to an empty 96-well plate (conditioned media (CM) plate). AP reaction solution (10 mM *p*-nitrophenylphosphate (*p*-NPP), 120 mM Tris-HCl (pH 9.5), 40 mM NaCl, and 10 mM MgCl₂) was dispensed into the cell plates and the CM plates (80 μ l). Absorbance at 405 nm (Abs₄₀₅) of the plates was measured, using a microplate reader (SpectraMax 340 PC384, Molecular Devices), before and after 1-h or 2-h incubation at room temperature. Ligand-induced AP-TGF- α release was calculated as described previously. Unless otherwise noted, spontaneous AP-TGF- α release signal, which varies from 8%–30% of total AP-TGF- α expression depending on transfected conditions, was subtracted from ligand-induced AP-TGF- α release signal. Using the Prism 7 software (GraphPad Prism), the AP-TGF- α release signals were fitted to a four-parameter sigmoidal concentration-response curve, from which EC₅₀ and E_{max} values were obtained. GPCRs and ligands used in this study is listed in [Table S1A](#).

Calculation of G-protein-coupling score

We used a factor known as the relative intrinsic activity (RAi) (Ehlert et al., 1999) to calculate scores for G-protein coupling. For each sigmoidal curve of chimeric G α -expressed condition, we divided a maximal response (E_{max}) by a potency (EC₅₀) and normalized an E_{max}/EC_{50} value to a maximum value among 11 chimeric G α curves. The resulting dimensionless, relative E_{max}/EC_{50} (defined as RAi) parameter was then logarithmically (base 10) transformed to give a LogRAi value used to quantify coupling. To minimize the occurrence of outliers arising from experimental variations, especially for weak AP-TGF- α release signal, we set two thresholds. As a first threshold, a G α chimera condition in which E_{max} was smaller than 3% AP-TGF- α release or a concentration-response curve did not converge, was regarded as RAi value of 0. As a second threshold, RAi value smaller than 0.01 was set as 0.01. Thus, a LogRAi values range from -2 to 0 and for the bioinformatics analyses, we used mean values of LogRAi ($n = 3-6$; for details of each GPCRs, see [Table S1A](#) and [Data S1](#)).

NanoBiT-G-protein dissociation assay

Plasmid transfection was performed in a 6-well plate with a mixture of 100 ng G α -Lg-encoding plasmid, 500 ng Sm-G β -encoding plasmid, 500 ng untagged G γ ₂-encoding plasmid, 200 ng GPCR-encoding plasmid with or without 100 ng RIC8-encoding plasmid (per well, hereafter). Unless otherwise stated, the combination of following plasmid mixtures was used: G α _s-Lg, Sm-G β ₁, G γ ₂ and RIC8B for NanoBiT-G_s; G α _{i1}-Lg, Sm-G β ₅ and G γ ₂ for NanoBiT-G_{i1}; G α _{i2}-Lg, Sm-G β ₃ and G γ ₂ for NanoBiT-G_{i2}; G α _{i3}-Lg, Sm-G β ₃ and G γ ₂ for NanoBiT-G_{i3}; G α _o-Lg, Sm-G β ₁ and G γ ₂ for NanoBiT-G_o; G α _q-Lg, Sm-G β ₁, G γ ₂ and RIC8A for NanoBiT-G_q; G α ₁₂-Lg, Sm-G β ₁, G γ ₂ and RIC8A for NanoBiT-G₁₂; G α ₁₃-Lg, Sm-G β ₁, G γ ₂ and RIC8A for NanoBiT-G₁₃. For G₁₂-coupled DREADD screening experiment, the combination of following plasmid mixture was used: G α ₁₂-Lg, G β ₁, Sm-G γ ₁₁ and RIC8A for NanoBiT-G₁₂; G α _o-Lg, G β ₁ and Sm-G γ ₁₁ for NanoBiT-G_o. After 1-day culture, the transfected cells were harvested with 1 mL of 0.53 mM EDTA-containing Dulbecco's PBS (D-PBS), followed by addition of 2 mL the HEPES-containing HBSS. The cells were pelleted by centrifugation at 190 g for 5 min and resuspended in 2 mL of the 0.01% BSA- and 5 mM HEPES (pH 7.4)-containing HBSS (assay buffer). The cell suspension was seeded in a 96-well culture white plate (Greiner Bio-One) at a volume of 80 μ l (per well hereafter) and loaded with 20 μ l of 50 μ M coelenterazine (Carbosynth) solution diluted in the assay buffer. After 2-h incubation with coelenterazine at room temperature, background luminescent signals were measured using a luminescent microplate reader (SpectraMax L, Molecular Devices). We note that incubation time with coelenterazine can be shortened, but an effect of baseline drift should be taken into account ([Figures S6C–S6E](#)). Test compound (6X, diluted in the assay buffer) was manually added to the cells (20 μ l). Luminescent signals were measured 3-5 min after ligand addition and divided by the initial count. The ligand-induced signal ratio was normalized to that treated with vehicle. The consequent fold-change values were fitted to a four-parameter sigmoidal concentration-response described above.

NanoBiT-RhoA assay

Plasmid transfection in HEK293 cells was performed by using a mixture of 100 ng Lg-RhoA plasmid, 500 ng Sm-PKN1 plasmid and 200 ng GPCR plasmid (per well in a 6-well plate). For transfection in MDA-MB-231 cells and PC-3 cells, 1.5 μ g Lg-RhoA plasmid and 7.5 μ g of Sm-PKN1 plasmid were used (per 10-cm dish). The transfected cells were harvested, seeded in a white 96-well plate and loaded with 10 μ M CTZ in the same manner described in the NanoBiT-G-protein dissociation assay. After measuring an initial luminescent signal, test compounds were added to the cells. Then, 3–5 min later, luminescent signals were measured and fold-change values were plotted as described above.

NanoBiT-IP₃ sensor assay

Plasmid transfection was performed by using a mixture of 1 μ g Lg-IP3R2-Sm plasmid and 200 ng GPCR plasmid (per well in a 6-well plate). The transfected cells were harvested, seeded in a white 96-well plate and loaded with 10 μ M CTZ in the same manner described in the NanoBiT-G-protein dissociation assay. After measuring an initial luminescent signal, test compounds were added to the cells. Then, 5–10 min later, luminescent signals were measured and fold-change values were plotted as described above.

siRNA transfection

Stealth siRNA duplexes against mRNA encoding $G\alpha_q$, $G\alpha_{11}$, $G\alpha_{12}$, $G\alpha_{13}$ and TACE (gene symbols, *GNAQ*, *GNA11*, *GNA12*, *GNA13* and *ADAM17*, respectively) and Stealth negative control were purchased from Life Technologies. Target sequences and manufacturer's catalog numbers are as follows: *GNAQ* (#1), 5'-GGAGAGAGUGGCAAGAGUACGUUUA-3', GNAQHSS104236; *GNAQ* (#2), 5'-CCCUUUGACUUACAAGUGUCAUUU-3', GNAQHSS104237; *GNA11* (#1), 5'-CCGGCAUCAUCGAGUACCCUUUCGA-3', GNA11HSS178464; *GNA11* (#2), 5'-GCAUCAGUACGUCAGUGCCAUCAAG-3', GNA11HSS104213; *GNA12* (#1), 5'-CCAAGGAAUUGUGGAGCAUGACUU-3', GNA12-HSS178466; *GNA12* (#2), 5'-CCAUCGUCAACAACAAGCUCUUCUU-3', GNA12MSS204749; *GNA13* (#1), 5'-CAGAAGCCCUUAUACCACCACUUCA-3', GNA13-HSS173827; *GNA13* (#2), 5'-GCAGCCCAA GGAAUGGUGGAAACAA-3', GNA13-HSS116479; *ADAM17*, 5'-CAGAAUCGUGUUGACAGCAAAGAAA-3', ADAM17-HSS186181. siRNA constructs for the *GNA12* (#1), the *GNA13* (#1) and the *ADAM17* genes were described previously and validated (Inoue et al., 2012).

HEK293 cells were seeded in a 6-well culture plate at cell density of 1×10^5 cells ml^{-1} in 2 mL of the complete DMEM and incubated for 1 day. Transfection of siRNA transfection was performed by using Lipofectamine® RNAiMAX (Thermo Fisher Scientific) according to the manufacturer instructions (final siRNA concentration of 10 nM and 2 μ L (per well in a 6-well plate) of Lipofectamine® RNAiMAX). After 1-day incubation, media were replaced and transfection of plasmids encoding AP-TGF- α and GPCR was performed as described above. The resulting cells were subjected to the TGF- α shedding assay.

Quantitative real-time PCR analysis

Total RNA from siRNA-transfected HEK293 cells was prepared using a GenElute Mammalian Total RNA Miniprep Kit (Sigma-Aldrich). Total RNA was reverse-transcribed using High-Capacity cDNA RT Kits (Applied Biosystems) according to manufacturer instructions. Real-time quantitative PCRs were performed with SYBR Premix Ex Taq (Takara Bio) and monitored by ABI Prism 7300 (Applied Biosystems). Standard plasmids ranging from 10^2 – 10^8 copies per well were used to quantify the absolute number of transcripts of cDNA samples. The numbers of transcripts were normalized to the number of *GAPDH* in the same sample and expressed as relative values to that in control siRNA-transfected cells. Primers were as follows: *GNAQ*, 5'-ACCGAATGGAGGAAAGCAAGG-3' and 5'-CATCTCTCTGGGGTCCATCATATTC-3'; *GNA11*, 5'-CAGCGAATACGACCAAGTCC-3' and 5'-ACCAGGGGTAGGTGATGATG-3'; *GNA12*, 5'-GAGGGATTCTGGCATCAGG-3' and 5'-CGATCCGGTCCAAGTTGTC-3'; *GNA13*, 5'-CCTGGATAACTTGGATAAACTTGG-3' and 5'-TTCATGGATGCCTTTGGTG-3'; *GAPDH*, 5'-GCCAAGGTCATCCATGACAAC-3' and 5'-GAGGGGCCATCCACAGTCTT-3'.

Western blot

The parental HEK293 cells and a panel of the G-protein-KO HEK293 cells (ΔG_q , ΔG_{12} and $\Delta G_q/\Delta G_{12}$ cells) in growth phase were harvested and approximately 1×10^6 cells were lysed in 500 μ L of SDS-PAGE sample buffer (62.5 mM Tris-HCl (pH 6.8), 50 mM dithiothreitol, 2% SDS, 10% glycerol and 4 M urea) containing 1 mM EDTA and 1 mM phenylmethylsulfonyl fluoride. Cell lysates were homogenized with a hand-held ultrasonic homogenizer (Microtech) and proteins were denatured at 95°C for 5 min. The lysates were loaded and separated on a 12.5% polyacrylamide SDS-gel. After electrophoresis, the gel was blotted to a nitrocellulose membrane. The blotted membrane was blocked with 5% skim milk-containing blotting buffer (10 mM Tris-HCl (pH 7.4), 190 mM NaCl and 0.05% Tween 20), immunoblot with primary (1 μ g ml^{-1}) and secondary antibodies (1:2000 dilution). Primary antibodies used in this study were anti- $G\alpha_q$ antibody (goat polyclonal; Abcam, ab128060), anti- $G\alpha_{11}$ antibody (mouse monoclonal, clone D-6; Santa Cruz Biotechnologies, sc-390382), anti- $G\alpha_{13}$ antibody (rabbit monoclonal, clone EPR5436; Abcam, ab128900) and anti- α -tubulin antibody (mouse monoclonal, clone DM1A; Santa Cruz Biotechnologies, sc-32293). We note that by using cell lysates overexpressing $G\alpha$ subunits, the anti- $G\alpha_q$ antibody and the anti- $G\alpha_{13}$ antibody were validated to be specific, but the anti- $G\alpha_{11}$ antibody reacted with both $G\alpha_q$ and $G\alpha_{11}$ (data not shown), and thus we labeled immuno-reactive bands as $G\alpha_{q/11}$. Secondary antibodies were conjugated with horseradish peroxidase (HRP) and were anti-goat IgG antibody (American Qualex, A201PS), anti-mouse IgG (GE Healthcare, NA9310) and anti-rabbit IgG (GE Healthcare, NA9340). Membrane were soaked with a commercial chemiluminescent

reagent (ImmunoStar® Zeta, FujiFilm Wako Pure Chemicals) or in-house reagent (100 mM Tris-HCl (pH 8.5), 50 mg ml⁻¹ Luminol Sodium Salt HG (FujiFilm Wako Pure Chemicals), 0.2 mM *p*-Coumaric acid and 0.03% (v/v) of H₂O₂), and a chemiluminescence image was acquired with a LAS-4000 (FujiFilm) and analyzed with Multi Gauge ver. 3.0 (FujiFilm).

Flow cytometry

Plasmid transfection was performed in a 12-well plate with volumes of 500 ng plasmid encoding N-terminally FLAG epitope-tagged GPCR with or without 250 ng *G α* -encoding plasmid. The transfected cells were harvested by adding 300 μ l of 0.53 mM EDTA-containing D-PBS, followed by 300 μ l of 5 mM HEPES (pH 7.4)-containing Hank's Balanced Salt Solution (HBSS). The cell suspension was dispensed in a 96-well V-bottom plate (200 μ l per well, two wells per sample). After centrifugation at 700 g for 1 min, the cells were washed once with D-PBS and pelleted. Cell pellets were suspended in 2% goat serum- and 2 mM EDTA-containing D-PBS (blocking buffer; 100 μ l per well) and incubated for 30 min on ice. After centrifugation at 700 g for 1 min, the cells were stained with anti-FLAG epitope tag monoclonal antibody (Clone 1E6, FujiFilm Wako Pure Chemicals; 10 μ g ml⁻¹ in the blocking buffer; 50 μ l per well) for 30 min on ice. After rinse with D-PBS, cells were labeled with a goat anti-mouse IgG secondary antibody conjugated with Alexa Fluor 488 (Thermo Fisher Scientific; 10 μ g ml⁻¹ dilution in the blocking buffer; 25 μ l per well) for 15 min on ice. The cells were washed once with D-PBS, resuspended in 100 μ l of 2 mM EDTA-containing-D-PBS and filtered through a 40 μ m filter. The fluorescently labeled cells (approximately 20,000 cells per sample) were analyzed by an EC800 flow cytometer (Sony). Fluorescent signal derived from Alexa Fluor 488 was recorded in an FL1 channel and flow cytometry data were analyzed by a FlowJo software (FlowJo). Values of mean fluorescence intensity (MFI) were used for quantification.

GloSensor cAMP assay

Plasmid transfection was performed in a 6-well plate with a mixture of 1 μ g Glo-22F cAMP biosensor-encoding pCAGGS plasmid (gene synthesized with codon optimization by Genscript), 200 ng AVPR2-encoding plasmid and 100 ng of *G α_s* -Lg-encoding plasmid or native *G α_s* -encoding plasmid. After 1-day incubation, the transfected cells were harvested with 0.53 mM EDTA-containing D-PBS, centrifuged at 190 g for 5 min and suspended in 0.01% BSA- and 5 mM HEPES (pH 7.4)-containing HBSS (vehicle; 0.6 mL per well). The cells were seeded in a half-area white 96-well plate (Greiner Bio-one; 30 μ L per well) and loaded with D-luciferin potassium solution (10 μ L of 8 mM solution per well; FujiFilm Wako Pure Chemical, Japan). After 2 h incubation in the dark at room temperature, the plate was read for its initial luminescent count (integration time of 1 s per well; Spectramax L, Molecular Devices, Japan). The cells were treated with vehicle, arginine vasopressin (Peptide Institutes, Japan) or 10 μ M forskolin (FujiFilm Wako Pure Chemical, Japan) (10 μ L of 5X solution per well). Kinetics values were measured on the plates for 20 min and expressed as fold-change values. To obtain a concentration-response curve, fold-change luminescent signals at 10-min after compound addition were normalized to that in forskolin-treated condition. Using the Prism 7 software (GraphPad Prism), the cAMP signals were fitted to a four-parameter sigmoidal concentration-response curve, from which EC₅₀ values were obtained.

For the chimeric *G α_s* -based cAMP assay, ΔG_s cells were transfected with a mixture of 1 μ g Glo-22F plasmid, 200 ng GPCR plasmid and 100 ng chimeric *G α_s* plasmid containing the backbone of human *G α_s* subunit (short isoform, residues 1-374) and a substitution of C-terminal 6-amino acids. Amino acid sequences for the Glo-22F cAMP sensor and the chimeric *G α_s* subunits are listed in [Data S3](#). The transfected cells were harvested, seeded in the half-area 96-well plate, loaded with D-luciferin and stimulated with a GPCR ligand in the same manner as described above. Scores of G-protein-coupling values (RAi) were calculated as described in the TGF- α shedding assay section.

Active RhoA pulldown assay

HN12 cells and Cal27 cells were cultured to 50% confluency, and then serum starved overnight. To induce RhoA activation, cells were treated with 5 μ M LPA, 1 μ M Ang II, or 10 μ M CP-55940 for 5 min. Active RhoA levels were measured using the RhoA Pull-Down Activation Assay Biochem Kit (bead pull-down format; Cytoskeleton) following the manufacturer instruction using a modified lysis buffer (50 mM Tris-HCl (pH 7.2), 500 mM NaCl, 10 mM MgCl₂, 0.1% SDS, 1% NP-40). Briefly, after stimulation, samples were lysed and protein concentrations were quantified using DC Protein Assay (BioRad). Samples were adjusted to the same concentration with lysis buffer and 500 μ g of each protein lysate was added to 15 μ L GST-tagged Rhotekin-RBD bound to Sepharose beads. Samples were incubated while rocking at 4°C for 1.5 h. Beads were then washed, eluted in Laemmli sample buffer, and analyzed by western blot using a mouse monoclonal anti-RhoA antibody (Cytoskeleton).

Ca²⁺ mobilization assay

Plasmid transfection was performed in the parental, ΔG_q and ΔG_{12} HEK293 cells by using GPCR-encoding plasmid (AGTR1 or EDNRA; 5 μ g per 10-cm culture dish). After one-day incubation, the transfected cells were harvested with trypsinization. After centrifugation, the cells were suspended in serum-free DMEM at a cell concentration of 5×10^5 cells ml⁻¹, and 40 μ l (per well hereafter) of the cell suspension seeded in a half-area, clear-bottom black plate. The cells were further incubated in the incubator for one day. After loading 40 μ l of a Ca²⁺ indicator (FLIPR Calcium 5 Assay Kit, Molecular Devices) according to manufacturer instructions in the presence of 2.5 mM probenecid for 1 h in the incubator, the cell plate was placed in a fluorescence microplate reader (FlexStation 3,

Molecular Devices). Fluorescent signal was measured with automated pipetting of test ligands (20 μ L of 5X compounds). Fluorescent signals from 40 to 55 s after ligand addition were averaged and normalized to an initial count and expressed as a relative value to vehicle treatment.

Comparison of data from the chimeric G-protein-based assay with known couplings

We performed Receiver Operating Characteristic (ROC) analysis to compare the chimeric G-protein-based TGF- α shedding assay results to primary or secondary couplings from GtoPdb (Harding et al., 2018), defined as binary classifiers. We defined the optimal LogRAi cutoff as that maximizing the True Positive Rate (TPR, or sensitivity) while minimizing the True Negative Rate (TNR, or 1-specificity). We defined positives as GtoPdb couplings reported in at least 3 references, and negatives as the couplings that were never reported for these more studied receptors. We obtained a value close to -1 as the optimal LogRAi cutoff considering all G proteins altogether (Figure S7; Table S1D), which we then considered as a lower and upper confidence bound for positively and negatively coupled receptors.

Sequence-based coupling-determinant features

We first generated a multiple sequence alignments (MSAs) of the 144 Class A GPCR sequences (Data S4) using HMMalign from the HMMer package (Eddy, 1998), using the 7tm_1 Pfam (Finn et al., 2016) Hidden Markov Model (HMM). As in a previously described procedure (Hannenhalli and Russell, 2000), we subdivided the pool of receptor sequences into positively and negatively coupled to a given G protein using the optimal LogRAi cutoff as a lower and upper bound. These sub-alignments were used to build corresponding HMM profiles through hmmbuild (<http://www.hmmer.org/>), leading to 22 HMMs (coupled versus uncoupled for 11 G proteins).

From coupled and uncoupled HMM profiles for each G protein, we then extracted alignment positions present in both HMM models and showing statistically different distributions (Wilcoxon's signed-rank test; p value ≤ 0.05) of the 20 amino acid bit scores (Figure 4A). We also considered those alignment positions with consensus columns (i.e., those having a fraction of residues, as opposed to gaps, equal or greater than the hmmbuild's *symfrac* parameter, using default value of 0.5) present in either of HMM models. In details, if a consensus column was present only in the HMM profile of either the coupled or uncoupled groups, we labeled it as *insertion* or *deletion*, respectively. As additional features, we also included length and amino acid composition of the N- and C-termini (N-term and C-term) and the extra- and intra-cellular loops (ECLs and ICLs). For every G protein, only statistically significant (p value < 0.05 ; Wilcoxon's rank-sum test) features were considered.

To identify each positions within the alignment, we employed the Ballesteros/Weinstein scheme (Weinstein, 1995), using the consensus secondary structure from the 7tm_1 HMM model to number residues within helices in a consecutive way (Table S3C). Most conserved positions within each helix were defined according to GPCRDB (<https://www.gpcrdb.org>) (Isberg et al., 2017). We adjusted the B/W numberings for TM6, which we started at position 6.25 (domain position 200) instead of 6.31 (domain position 206), according to visual inspections of recent G protein-GPCRs complexes. If a position lies on an extra-7TM region (e.g., ECLs or ICLs), we use the corresponding label plus the corresponding Pfam domain consecutive numbering in parenthesis.

G-protein-coupling predictor

We implemented a predictor for G-protein coupling by using a logistic regression classifier, or Log-reg classifier, available from the scikit-learn package (<https://scikit-learn.org>) (Pedregosa et al., 2011). The possible outcomes in log-reg are modeled using a logistic function, with L1 or L2 based regularization. In this study we used L2 penalized form of log-reg. The target value is expected to be a linear combination of the given features. This property of log-reg can also be exploited to study the weights of its features.

As an optimization problem, binary class L2 penalized logistic regression minimizes the following cost function:

$$\min_{w,c} \frac{1}{2} w^T w + C \sum_{i=1}^n \log(\exp(-y_i(X_i^T w + c)) + 1) \quad (1)$$

where \mathbf{X} denotes a vector of feature variables, $\mathbf{w} \in \mathbf{R}^n$ is the weight vector, $\mathbf{c} \in \mathbf{R}^n$ is the intercept, \mathbf{C} is inverse of regularization strength (positive float), y takes values in $\{-1, 1\}$ at trial i and n is the number of trials conducted.

We used the liblinear method as the optimization algorithm as shown to be optimal for relatively small datasets (<https://www.csie.ntu.edu.tw/~cjlin/liblinear/>).

Training and cross validation

We used 7TM domain positions and compositional features for the ICL3 and C-term, which prevail over other extra-7TM domain features, to create a training matrix. In case of significant positional features, two-bit scores (derived from the positive and negative HMMs for a given G protein) are returned for the corresponding amino acid found at a given position in the input GPCR sequence (Figure 4A). In case a position was found to be present in either positive or negative HMMs, the single bit score, derived from the respective HMM, was returned. If for any GPCR, no amino acid was present at the given position, it was assigned the highest bit scores from the both models, implying the least conserved scores.

All the features were scaled to the range $[0, 1]$. Feature scaling aids not only in converging the algorithm faster but also helps in assessing the feature relevance (Dou et al., 2012). A grid search was performed over a stratified 5-fold cross validation (CV) to select

the best value of C (inverse of the regularization strength) over a range of $[1e-02, 1e05]$. In a stratified 5-fold CV, the training matrix is divided randomly into 5 equal sub-matrices, preserving the ratio of positive (coupling) and negative (non-coupling) GPCRs. During each fold, one of the sub matrix is treated as the validation set and the remaining four as the training set.

We assessed the performance of our predictor using standard metrics (MCC, ACC, PRE, REC, SPE, AUC, F1M; [Table S2A](#)). The parameters showing the best Area Under the Curve (AUC) of the Receiver Operating Curve (ROC) were chosen to create models for every G protein ([Table S2A](#)).

The number of positive (coupling) GPCRs were either more (eg: in GNAI1/3) or less (eg: in GNAS) than the number of negative (non-coupling) GPCRs. Such an imbalance would make the predictor biased to any one of the two classes. In order to counter this problem, the parameter `class_weight` was set to *balanced* in the log-reg classifier function. By default, all the classes have same weight. However, by setting the `class_weight` as *balanced*, the values of the column with classes (coupling/uncoupling) are used to automatically adjust the weights inversely proportional to their frequencies in the training matrix. To ensure minimal variance due to random division of the training matrix during the cross validation, the aforementioned experiment was repeated ten times for every G-protein group and the standard deviation was recorded. The feature weights were extracted as described elsewhere ([Dou et al., 2012](#)) from the trained models and are critical to understand the relative importance of different features ([Figure 4](#)).

Besides performing the above-said steps at LogRAi cutoff of -1.0 , we also created models at LogRAi cutoffs -0.5 and -0.1 . As it can be seen in [Figure S7A](#), for most of the G proteins, -1.0 turns out to be the best LogRAi cutoff during cross-validation (using MCC as the selection criteria).

Randomized training test

In order to assess over-fitting, we performed a randomization test ([Salzberg, 1997](#)). For every G protein, the original labels of the training matrix were replaced with randomly determined labels, while preserving the ratio of number of positive (coupling) and negative (non-coupling) GPCRs. Performance using the randomization training set was lower than that of actual training set ([Table S2B](#)), implying that our strategy is insensitive to the data training set.

Test set to benchmark predictor performance

To benchmark our method and compare it with Pred Couple ([Sgourakis et al., 2005b](#)), a web-server available to predict GPCR-G-protein coupling, we extracted all the GPCRs from GtoPdb that are present in neither the chimeric G-protein-based TGF- α shedding assay nor in Pred Couple's training set, thus obtaining a list of 86 unseen GPCRs ([Data S4](#)). As mentioned above, one of the major limitations of GtoPdb is the absence of a definite true negative set, thus, the best measure to compare our predictor with that of Pred Couple is *recall*, also known as sensitivity or the true positive rate. Since both GtoPdb and Pred Couple provide coupling information at the G-protein family level, we combined the performance of individual G-protein predictors based on their families to compare the performance of our method with Pred-Couple. For example: if a given GPCR was predicted to couple to at least one of the G proteins of a family, it was annotated as coupling to that G-protein family. The combinations of our predictors at the family level outperformed Pred-Couple over the test set ([Table S2C](#)). The individual G-protein predictors' and their combined (G-protein family level) performance over the Test set at different LogRAi cutoffs are reported in [Tables S2A](#) and [S2B](#) and [Figures S7A](#) and [S7C](#), respectively.

To further check the predictor's performance, we trained and tested an additional predictor using exactly the same procedure as reported above using GtoPdb coupling information instead of the TGF- α shedding assay ([Figure 4A](#); [Table S2D](#)).

Functional classification of coupling features through 3D structure analysis

We identified functional positions as those mediating inter- and intra-molecular contacts, i.e., whenever at least one pair of atoms, from either a residue-residue or residue-ligand interface, was found spatially closer than 5\AA . We analyzed 246 3D structures, representing 51 members of the GPCR Class A (PFAM: 7tm_1) family using PDB-Swissprot-PFAM correspondences available from SIFT (as of July 2018) ([Velankar et al., 2013](#)).

To define GPCR-ligand contact sites, we restricted our analysis only to GPCR putative ligands as defined in GtoPdb ([Harding et al., 2018](#)). We performed similarity searches between GtoPdb and PDB ligands using topological fingerprints from RDKit (<http://www.rdkit.org/>) generated from SMILES descriptors and we considered only the best matching GtoPdb ligand for a given PDB component. All the protein residues mediating contacts were mapped to protein sequence position using alignments between Uniprot canonical sequences and corresponding PDB generated through Blast ([Altschul et al., 1990](#)). Note that through this procedure we considered contacts mediated by the equivalent residues from different structures only once, thus avoiding overcounting due to PDB redundancy. We then mapped the amino acids found in contact with putative ligands on the PFAM multiple sequence alignments (MSA). Based on available GPCR-G-protein complexes (PDB: 3SN6, 5G53, 6D9H, 6GDG, 6DDE, 6DDF, 6CMO and 6D9H) we similarly identified the residues forming the receptor-G-protein interaction interface by using a distance cutoff of 6.5\AA to define atom-pairs forming inter-residue contacts.

Similarly to methods employed to decipher the activation mechanisms of GPCRs ([Angelova et al., 2011](#); [Venkatakrishnan et al., 2016](#); [Venkatakrishnan et al., 2013](#)) and other signaling molecules ([Papaleo et al., 2016](#); [Raimondi et al., 2013a](#); [Raimondi et al., 2013b](#); [Seeber et al., 2015](#)), we also inspected the network of intramolecular contacts using the same thresholds as above and we similarly mapped the identified positions on Class A 7TM MSA. We then defined a consensus contact network by considering the sequence positions (nodes) found in contact (edges) in at least 50% of the analyzed sequences. We performed network analysis

through *igraph* (Csardi and Nepusz, 2006), defining as hub nodes having a degree of at least 4. We generated functional state consensus networks by grouping available structures using ligand classification from GtoPdb (i.e., agonist or antagonist/inverse agonist) or functional classification directly available from the protein databank (i.e., active or inactive), thus defining *active-like* (i.e., agonist-bound/active) or *inactive-like* (i.e., antagonist-bound/inactive) states (Table S3D). Structures where this classification was not possible were discarded.

We calculated the shortest paths connecting positions forming the consensus ligand and G-protein-binding interfaces within active- and inactive-like networks through the Dijkstra algorithm (Dijkstra, 1959) from *igraph*. Active-state specific shortest paths were defined as those characterized by having either endpoints, or intermediate connectivity residues, exclusive to active-like state contact network.

Predictions of G₁₂-coupled DREADD chimeric sequences

In order to predict mutant sequences with enhanced G_{12/13} coupling capabilities, we started from the available DREADD coupled with G_{q/11} (M3D) and G_{12/13} (M4D) (Armbruster et al., 2007). We generated chimeric sequences by swapping on these backbones the ICL3 and C-term sequence stretches derived from each receptor of the chimeric G-protein-based assay panel (148 GPCRs). We first aligned the receptor sequences, including M3D and M4D, to the PFAM 7tm_1 HMM model. We defined ICL3 as the MSA region comprised within HMM positions 173-205, and the C-term as the MSA portion starting after 7tm_1 HMM end (i.e., position 268). We then created M3D and M4D chimeras by exchanging their ICL3 and C-term sequences with the corresponding sequences from each receptor tested in the chimeric G-protein-based TGF- α shedding assay. We generated 296 chimeric sequences by swapping the ICL3 alone or in combination with the C-termini.

We then predicted the coupling probability to GNA12/GNA13 for each chimeric sequence, ranking them according to their relative coupling probability (i.e., $\Delta\text{Pred_Coup} = \text{Pred_Coup}^{\text{DREADD_MUT}} - \text{Pred_Coup}^{\text{DREADD}}$). We selected the top 10 chimeric sequences for experimental validation.

QUANTIFICATION AND STATISTICAL ANALYSIS


Statistical analyses were performed using GraphPad Prism 7 software and methods are described in the legends of the figures. In flow cytometry experiments, approximately 20,000 cells were measured for their fluorescent signals and data were analyzed by FlowJo software. Mean fluorescent intensity was used for quantification of cell surface GPCR expression. Representation of symbols and error bars is described in the ligands. Symbols are either mean values of indicated numbers of independent experiments or data-point from single experiment. Error bars denote SEM or SD. Concentration-response curves were fitted to all data by the Nonlinear Regression: Variable slope (four parameter) in the Prism 7 tool. Linear regression and representation of 90% confidence bands were performed by the Prism 7 tool. For multiple comparison analysis in the flow cytometry data and G₁₂-DREADD generation, two-way ANOVA and following Dunnett's test and Sidak's test, respectively, was used.

DATA AND SOFTWARE AVAILABILITY

The Python code used for the predictor is available on GitHub (https://github.com/raimondifranc/gpcr_coupling_predictor). Heat-maps in Figure 2 and 5 were generated using scipy (<https://www.scipy.org/>) and matplotlib python libraries (<https://matplotlib.org/>).

ADDITIONAL RESOURCES

We have created an accessible resource for interrogating the data generated by the G-protein-based assay and a tool for online predictions of coupling based on sequences available at <http://gpcr.russelllab.org/>. The GPCR/G-protein interactions with a score LogRAi ≥ -1 have been annotated and are accessible in the SIGNOR database (<https://signor.uniroma2.it/downloads.php>).



(legend on next page)

Figure S1. Validation of the TGF- α Shedding Assay, Related to Figures 1, 2, and 3

(A) siRNA-mediated knockdown of mRNA expression. HEK293 cells transfected with a siRNA construct specific to each gene (two targeting constructs per gene) were analyzed for mRNA expression by quantitative real-time PCR. The *GNAQ*, the *GNA11*, the *GNA12* and the *GNA13* genes encode G_{α_q} , $G_{\alpha_{11}}$, $G_{\alpha_{12}}$, and $G_{\alpha_{13}}$ subunits, respectively. mRNA levels are shown as relative values to that in control siRNA-transfected cells. Bars and error bars represent mean and SEM, respectively ($n = 3$).

(B) siRNA-mediated knockdown at protein expression levels. Lysates from HEK293 cells transfected with a mixture of the indicated siRNA constructs were subjected to immunoblot analyses using antibodies specific to G_{α_q} (an open arrowhead), dually to G_{α_q} and $G_{\alpha_{11}}$ ($G_{\alpha_q/11}$), $G_{\alpha_{13}}$ or α -tubulin. Note that owing to a lack of a sensitive, validated antibody against $G_{\alpha_{12}}$, immunoblot for $G_{\alpha_{12}}$ was not assessed. ns, non-specific immunoreactive band (a filled arrowhead).

(C) Knockdown of $G_{\alpha_{11}}$ attenuates AP-TGF- α release induced by $G_{\alpha_{11}}$ -coupled HRH1. HEK293 cells transfected with a siRNA construct (filled symbols) and an HRH1-encoding plasmid were subjected to the TGF- α shedding assay. Note that the data for the control siRNA (open symbols) are identical in all of the panels. The *ADAM17* gene encodes a membrane protease that cleaves the AP-TGF- α reporter protein (Inoue et al., 2012). Numbers to the right of each plot indicate EC_{50} , E_{max} and RA_i values obtained from sigmoidal concentration-response curves. Symbols and error bars are mean and SD (three replicate wells per one point), respectively, from a representative experiment of at least two independent experiments with similar results.

(D) Knockdown of $G_{12/13}$ attenuates AP-TGF- α release induced by $G_{12/13}$ -coupled PTGER3. Details as for (C), but using another PTGER3 and the corresponding ligand, prostaglandin E_2 (PGE_2).


(E) Parental, ΔG_q , ΔG_{12} and $\Delta G_q/\Delta G_{12}$ HEK293 cells were transiently transfected with a plasmid encoding N-terminally FLAG epitope-tagged GPCR (HRH1, ADRB1 or DRD1) or an empty plasmid (Mock). The transfected cells were stained fluorescently labeled with anti-FLAG tag antibody, followed by a secondary antibody conjugated with a fluorophore, and subjected to flow cytometry analysis. Numbers at bottom of each histogram indicate mean \pm SD (four biological replicates) of percentage of receptor-positive cells (a bar region in the histogram) and mean fluorescent intensity (MFI) from a representative of two independent experiments with similar results. Note that due to transient transfection, there are two peaks showing a highly expressing cell pool and poorly expressing one.

(F) In $\Delta G_q/\Delta G_{12}$ HEK293 cells, chimeric G_{α} subunits were individually transfected with a plasmid encoding an N-terminally FLAG epitope-tagged GPCR (HRH1, ADRB1 or DRD1). The transfected cells were stained fluorescently labeled with anti-FLAG tag antibody, followed by a secondary antibody conjugated with a fluorophore, and subjected to flow cytometry analysis. Plots in the panels denote independent experiments and lines represent mean values ($n = 4$ or 5). MFI values that significantly differ from the control (G_{α_q} (ΔC)) are denoted by asterisks: * $p < 0.05$, ** $p < 0.01$ (one-way ANOVA with Dunnett's post hoc test). NS denotes not significantly different from the control.

(G) Concentration-response curves of AGTR1 for the endogenous ligand (Angiotensin II, AngII) and a biased agonist ($[Sar^1-Ile^4-Ile^8]$ AngII, SII). G-protein signaling activity was assessed by the chimeric-G-protein-based assay. Symbols and error bars are mean and SEM, respectively, of eight independent experiments with each performed triplicate.

(H) Ligand bias plots. For each chimeric-G-protein coupling, LogRA_i values were plotted. If SII behaves as a balanced ligand, plots would be linearly aligned. Dotted lines (slope = 1) were drawn crossing C-terminal G_{α_q} or $G_{\alpha_{12}}$ chimera, indicating that SII is more biased toward G_{12} than G_q . Note that activation of G_{α_s} by SII was minimum and thus not included in the plot. Symbols and error bars are mean and SEM, respectively, of eight independent experiments.

(I) Validation of SII bias toward G_{12} by the NanoBIT-G-protein dissociation assay. NanoBIT-G-proteins (G_q and G_{12}) were expressed with AGTR1 and ligand-induced G-protein-dissociation signal was measured. Symbols and error bars are mean and SEM, respectively, of six independent experiments.



(legend on next page)

Figure S2. Chimeric G α Subunits and Their Activity for TGF- α Shedding and cAMP Responses, Related to Figures 1, 2, and 3

(A) Seven C-terminal sequences (CGN numbering (Flock et al., 2015) of G.H5.20-G.H5.26) of G α subunits among the 16 human G α subunits. Asterisks indicate identical amino acids to one above. Note that there are 11 distinct sequences and that the -7 position (G.H5.20) is a completely conserved leucine.

(B) Evolutionary conservation the seven C-terminal sequences of representative G α subunits from the four G-protein subfamilies.

(C) Chimeric G-proteins used in this study. We used human G α_q -based chimera with a substitution of six C-terminal amino acids. In the negative-control G α_q , the seven C-terminal amino acids are truncated.

(D) Capacity of G α subunits to induce TGF- α shedding response. Scheme of the experiment is shown in left. G α_o -coupled DRD2 was co-expressed with an indicated chimeric G α subunit or a native, full-length G α subunit in $\Delta G_q/\Delta G_{12}$ cells and subjected to the TGF- α shedding assay by using dopamine. Note that the G $\alpha_{q/11}$ chimera induced the most potent response and the other negative control conditions (G α_{11} , G α_q or an empty vector transfection (Mock)) did not induce the signal. Symbols and error bars are mean and SD (three wells per one point), respectively, from a representative experiment of at least two independent experiments with similar results.

(E) Capacity of G α subunits to induce TGF- α shedding response. Scheme of the experiment is shown in left. The experimental design is the same as DRD2, except for usage of G α_s -coupled PTGER2, C-terminal G α_s chimeras and prostaglandin E $_2$ (PGE $_2$). Note that the G $\alpha_{q/s}$ chimera induced the most potent response and the other negative control conditions (G α_s long isoform (G α_{sL}), G α_s short isoform (G α_{sS}), G α_q or an empty vector transfection (Mock)) did not induce the signal. Symbols and error bars are mean and SD (three replicate wells per one point), respectively, from a representative experiment of at least two independent experiments with similar results.

(F) Protein expression levels of chimeric G α_q subunits. Lysates from $\Delta G_q/\Delta G_{12}$ cells transfected with a plasmid encoding an indicated chimeric G α_q subunit were subjected to immunoblot analyses using antibodies specific to G α_q , G $\alpha_{q/11}$ or α -tubulin. Note that expression levels of the chimeric G α_q subunits were almost equal except for the C-terminally truncated G α_q (ΔC), which was previously shown to undergo spontaneous activation (Denker et al., 1992), and thereby likely to be unstable in cells owing to its tendency to separate from G $\beta\gamma$ subunits.





(G) Kinetics of cAMP level upon G α_s -coupled receptor stimulation. HEK293 cells devoid of the G α_s subfamily (ΔG_s , lacking G α_s and G α_{oit}) (Stallaert et al., 2017) transiently expressing a cAMP biosensor (Glo-22F), a G α_s -coupled receptor (AVPR2 or mock transfection) and a G α_s construct (native G α_s , G α_s -Lg or mock transfection) were loaded with D-luciferin. The cells were stimulated with an increasing concentration of arginine-vasopressin, an AVPR2 ligand, or Forskolin (FSK, 10 μ M), an adenylyl cyclase activator, and luminescent signals were measured for 20 min. Luminescent signals were normalized to initial counts and relative values are plotted. Each line indicates a kinetics from a single well and data are from a representative experiment of at least three independent experiments with similar results. Note that owing to preference of Forskolin to a G α_s -bound adenylyl cyclase (Insel and Ostrom, 2003), Forskolin-induced cAMP response was attenuated in the ΔG_s cells. Also note that in native G α_s -expressing cells, owing to higher initial luminescent counts reflecting constitutive G α_s activity, amplitude of fold change is smaller than the other conditions.

(H) Concentration-response curves. Fold-change luminescent signals at 10 min after ligand addition in A were normalized to Forskolin response and fitted to a sigmoidal curve. Symbols and error bars are mean and SEM, respectively (n = 3 or 4). Pharmacological parameters are shown at the bottom (mean \pm SEM). Mean pEC $_{50}$ values were anti-logarithmically transformed and expressed as pM values in parenthesis.

(I) Chimeric G-protein-based cAMP assay in ΔG_s cells. A test GPCR is expressed together with one of 11 chimeric G α_s subunits harboring C-terminal 6-amino acid substitution in ΔG_s cells and restoration of ligand-induced cAMP response is measured by a luminescent cAMP biosensor. The C-terminally truncated G α_s construct (G α_s (ΔC)) is used for a negative control.



(J) Representative data for the chimeric G-protein-based cAMP assay. TBXA2R was expressed with one of the 11 G α_s constructs or the G α_s (ΔC) and treated with titrated concentration of a ligand (U-46619). Ligand-induced cAMP responses normalized to forskolin (10 μ M)-induced response were fitted to a sigmoidal concentration-response curve (upper panels). G-protein coupling is scored as logarithm of RAI values. Symbol size is proportional to E_{max} , which reflects fitting quality. Data for the concentration-response curves are from a representative experiment (mean \pm SD of triplicate measurements). Each LogRAI plot denotes single experiment and bars and error bars represent mean and SEM, respectively (n = 5).

(K) Comparison of the chimeric G-protein backbones in coupling profiles. LogRAI values obtained from the chimeric G α_q -based TGF- α shedding assay are plotted against the chimeric G α_s -based cAMP assay for seven prostanoid receptors. Considered were only mean values for the plots. Note that PTGFR showed poor responses in the cAMP assay and thus not used for the comparison. Linear regression analysis was performed and 90% confidence bands of the best-fit line were shown. Mean \pm SD of r^2 values from the seven prostanoid receptors is shown at the bottom.



E

Pre-incubation with CTZ (min)	Δ Signal	pEC50	n	Δ Signal	pEC50	n	Δ Signal	pEC50	n	Δ Signal	pEC50	n
15	0.28 ± 0.01	8.56 ± 0.13	5	0.33 ± 0.03	7.72 ± 0.18	4	0.22 ± 0.02	8.49 ± 0.09	5	0.43 ± 0.01	9.50 ± 0.02	3
30	0.29 ± 0.01	8.41 ± 0.19	5	0.30 ± 0.01	7.92 ± 0.14	4	0.21 ± 0.01	8.51 ± 0.03	5	0.44 ± 0.01	9.37 ± 0.06	3
60	0.29 ± 0.01	8.65 ± 0.14	5	0.29 ± 0.02	7.92 ± 0.07	4	0.19 ± 0.02	8.64 ± 0.03	5	0.49 ± 0.02	9.38 ± 0.03	3
120	0.30 ± 0.01	8.70 ± 0.11	5	0.27 ± 0.01	7.83 ± 0.08	4	0.20 ± 0.01	8.42 ± 0.05	5	0.50 ± 0.02	9.35 ± 0.05	4



(legend on next page)

Figure S3. Development and Validation of the NanoBiT-G-Protein Dissociation Assay, Related to Figure 1

(A) Scheme of the NanoBiT-G-protein assay. Components of the NanoBiT-G-protein (typically, LgBiT-inserted $G\alpha$ subunit, SmBiT-fused $G\beta_1$ subunit and native $G\gamma_2$ subunit) and a test GPCR are transiently expressed in HEK293 cells. By loading with coelenterazine (CTZ), a substrate for the NanoBiT luciferase, the NanoBiT-G-protein emits bioluminescence. Stimulation with a GPCR ligand triggers exchange of a guanine nucleotide (GDP release and GTP incorporation) of the NanoBiT-G-protein and induces dissociation of $G\alpha$ -Lg from Sm- $G\beta\gamma$, thereby reducing bioluminescence signals. Note that both $G\alpha$ and $G\beta\gamma$ subunits are lipidated (not shown) and localized to membrane.

(B) Luminescent kinetics of NanoBiT-G-proteins after GPCR ligand stimulation. Representative members (G_s , G_{i3} , G_q and G_{13}) from the four G-protein subfamilies were co-expressed with a corresponding coupling GPCR and stimulated with its ligand (10 μ M isoproterenol, 10 μ M dopamine or 1 μ M U-46119) or vehicle. After ligand addition, a microplate was measured at 10 s interval for 10 min. Each line indicates a luminescent trace of one well, normalized to an initial count, from a representative experiment of at least three independent experiments with similar results. Note that due to a time lag between manual ligand addition and beginning of measurement, NanoBiT- G_{13} shows already dissociated signal from the initial reading.

(C-E) Effect of preincubation time with CTZ. Cells expressing the indicated combination of the NanoBiT-G-protein and the test GPCR were loaded with CTZ for 15-120 min before measurements (baseline and ligand stimulation). Luminescent signals after 3-5 min ligand addition (10 μ M isoproterenol, 10 μ M dopamine or 1 μ M U-46119) or vehicle treatment were normalized to the baseline signals (C). Bars and error bars represent mean and SEM of indicated numbers of independent experiments in E. Changes in the luminescent signals across titrated ligand concentrations were further normalized to that of vehicle treatment and expressed as concentration-response curves (D). Symbols and error bars represent mean and SEM of indicated numbers of independent experiments in E. Parameters (signal changes and pEC_{50} values) were calculated from the sigmoidal curves (E). Data are expressed as mean \pm SEM of indicated numbers of independent experiments. Mean pEC_{50} values were anti-logarithmically transformed and expressed as nM values.

(F) Validation of the NanoBiT-G-proteins by using prostanoid receptors. Seven NanoBiT-G-protein (G_s , G_{i2} , G_{i3} , G_o , G_q , G_{12} and G_{13}) were profiled for the eight prostanoid receptors with their ligands shown in parentheses. Each dot represent data from independent experiments and lines and error bars indicate mean and SEM (n = 5-9).

(G) Comparison of coupling profiling between the chimeric G-protein-based TGF- α shedding assay and NanoBiT-G-protein assay. LogRAi values obtained from the chimeric G-protein-based assay are plotted against G-protein dissociation signals from the NanoBiT-G-protein assay for the eight prostanoid receptors. Considered were only mean values for the plots. The G_i family contains data for three members ($G_{\alpha q/11}$, $G_{\alpha q/13}$ and $G_{\alpha q/o}$ chimeras; NanoBiT- G_{i2} , NanoBiT- G_{i3} and NanoBiT- G_o , respectively) and the G_{12} family includes data for two members ($G_{\alpha q/12}$ and $G_{\alpha q/13}$ chimeras; NanoBiT- G_{12} and NanoBiT- G_{13} , respectively). PTGER3- G_{12} data were excluded owing to the increased luminescent signal. Linear regression analysis was performed and 95% confident bands of the best-fit lines intervals were shown.




Figure S4. Analysis of the Chimeric G-Protein-Based Assay Dataset and Comparison with GtoPdb, Related to Figures 2 and 3

(A) Roc curve comparing the chimeric G-protein-based TGF- α shedding assay couplings with GtoPdb couplings: roc curves (<https://www.scikit-learn.org/>) were generated considering only GtoPdb best characterized primary couplings (i.e., reported in at least 3 publications) as binary classifier and TGF- α shedding assay values as scores. Roc curves were calculated either considering coupling values for all the G-proteins in the chimeric G-protein-based TGF- α shedding assay (gray) or by excluding poorly characterized couplings (i.e., GNA12, GNA13, GNA14, GNA15, GNAZ; red curve).

(B) Number of GPCRs coupled to G-proteins of the four families at different LogRAI thresholds in the chimeric G-protein-based TGF- α shedding assay as well as in GtoPdb.

(C) Distribution of the number of reported bindings (of any of the four G-protein families) for each receptor at different LogRAI thresholds in the chimeric G-protein-based TGF- α shedding assay as well as in GtoPdb.

(D-F) Fractions of specific couplings, i.e., receptors binding to members of only one G-protein family, in the chimeric G-protein-based TGF- α shedding assay (dark red and orange bars for LogRAI ≥ -0.1 and -1 couplings) and GtoPdb (black and gray bars for primary only and primary & secondary couplings); Venn diagrams with the numbers of receptors coupled to each G-protein family in the chimeric G-protein-based TGF- α shedding assay at higher LogRAI stringencies ≥ -0.5 (E) and ≥ -0.1 (F).

(G) Comparison of receptor sequence and coupling profile similarities: we calculated receptor pairwise sequence similarity by outputting distance matrices from *ClustalO* (Sievers et al., 2011). We compared receptor pairwise coupling similarity by calculating the distance matrix of coupling profiles (i.e., vectors containing LogRAI values for 11 G proteins) through the *pdist* function from *scipy* (<https://www.scipy.org/>).









Figure S5. Validation of RhoA Activation by the Newly Identified G_{12/13}-Coupled GPCRs, Related to Figure 3

(A) Scheme of the NanoBiT-RhoA sensor. The two fragments (LgBiT and SmBiT) of the NanoBiT luciferase are N-terminally fused to RhoA and its effector PKN1, respectively. Upon activation by exchanging GDP to GTP, GTP-bound RhoA interacts with PKN1, thereby increasing luminescent signals being measurable upon loading with CTZ.



(B) Luminescent kinetics of the NanoBiT-RhoA sensor after GPCR ligand stimulation. HEK293 cells expressing the sensor alone (Mock) or with a test GPCR (PTGER3) were stimulated with its ligand (prostaglandin E₂, PGE₂), lysophosphatidic acid (LPA) or vehicle. After ligand addition, a microplate was measured at 10 s interval for 10 min. Each line indicates a luminescent trace of one well, normalized to an initial count, from a representative experiment of at least three independent experiments with similar results. Note that LPA stimulates LPA receptors endogenously expressed in HEK293 cells.

(C) Validation of G_{12/13}-mediated signal of the NanoBiT-RhoA sensor. PTGER3 or LPAR6 was expressed with the NanoBiT-RhoA sensor in the parental, ΔG_q, ΔG₁₂ and ΔG_q/ΔG₁₂ HEK293 cells, and ligand-induced luminescent signals were measured. Symbols and error bars represent mean and SEM, respectively, of 4 (PTGER3) and 6 (LPAR6) independent experiments with each performed in duplicate.

(D) NanoBiT-RhoA activation by selected GPCRs. Test GPCRs including the newly identified G_{12/13}-coupled GPCRs (Figure 3E) were expressed together with the NanoBiT-RhoA sensor in HEK293 cells and ligand-stimulated luminescent signals were measured. Note that CP-55940-induced RhoA activation occurred in cells expressing CNR1, but not CNR2. Symbols and error bars represent mean and SEM, respectively, of 5-7 independent experiments with each performed in single measurement or duplicate.




(E) NanoBiT-RhoA activation through endogenously expressed GPCRs. PC-3 cells and MDA-MB-231 cells transiently expressing the NanoBiT-RhoA sensor alone were stimulated with vehicle, thrombin (1000 U L⁻¹) or LPA (1 μM), which is a potent inducer of RhoA activation in many cell types. Bars and error bars represent mean and SEM, respectively, of 3 independent experiments with each performed in triplicate. *p < 0.05, **p < 0.01 as compared with vehicle treatment (one-way ANOVA with Dunnett's post hoc test).

(F) RhoA pull-down assay to detect G_{12/13} activation by endogenously expressed GPCRs. HN12 cells and Cal27 cells were serum-starved and treated with 5 μM LPA, 10 μM CP-55940, 1 μM Ang II or vehicle. Cell lysates were subjected to the pull-down assay using Rhotekin-beads and precipitated GTP-bound RhoA proteins as well as input RhoA proteins (Total) were assessed by immunoblot analysis. Images of immunoblot membranes are representatives of two experiments with similar results.



B Chimeric G-protein-based TGF α shedding assay

	AGTR1		EDNRA			AGTR1		EDNRA	
	Parent	ΔG_{12}	Parent	ΔG_{12}		Parent	ΔG_{12}	Parent	ΔG_{12}
pEC50	9.48 \pm 0.06	9.54 \pm 0.05 ^t	9.71 \pm 0.08	9.56 \pm 0.02	pEC50	9.36 \pm 0.16	9.44 \pm 0.19	9.67 \pm 0.06	9.93 \pm 0.05
E_{max}	57.0 \pm 1.6	54.3 \pm 2.1	60.9 \pm 1.9	58.8 \pm 0.8	E_{max}	0.87 \pm 0.04	0.72 \pm 0.04	0.91 \pm 0.05	0.77 \pm 0.07
n	6	6	4	4	n	4	4	4	4



G NanoBiT-IP₃ assay




Figure S6. $G_{q/11}$ Signaling in the Absence of $G_{12/13}$ for GPCRs Coupled with $G_{q/11}$ and $G_{12/13}$, Related to Figure 3

(A) Protein expression levels of $G\alpha$ subunits. Lysates from the parent, ΔG_q , ΔG_{12} and $\Delta G_q/\Delta G_{12}$ cells were subjected to immunoblot analyses using antibodies against $G\alpha_q$ (an open arrowhead), $G\alpha_{q/11}$, $G\alpha_{13}$ or α -tubulin. Note that compensatory upregulation of $G\alpha$ subunits in ΔG_q cells (for $G\alpha_{13}$) or ΔG_{12} (for $G\alpha_q$ or $G\alpha_{11}$) was not observed. Also, note that owing to a lack of a sensitive, validated antibody against $G\alpha_{12}$, immunoblot for $G\alpha_{12}$ was not assessed. ns, non-specific immunoreactive band (a filled arrowhead).

(B) Parameters obtained from concentration-response curves of the chimeric G-protein-based TGF- α shedding assay (Figure 3E). Parameters are shown as mean \pm SEM of the indicated numbers of independent experiments.

(C) Ca^{2+} mobilization assay. The parent, ΔG_q and ΔG_{12} cells transiently expressing AGTR1 or EDNRA were loaded with a Ca^{2+} fluorescent dye and ligand-induced Ca^{2+} mobilization was assessed. Symbols and error bars represent mean and SEM of the indicated numbers of independent experiments with each performed in duplicate. Parameters (mean \pm SEM) obtained from the concentration-response curves are shown at the bottom.


(D) Scheme of the NanoBiT-IP₃ sensor. The two fragments (LgBiT and SmBiT) of the NanoBiT luciferase are N-terminally and C-terminally, respectively, fused to inositol-triphosphate (IP₃) receptor IP3R2. Upon activation of phospholipase C β and hydrolysis of phosphoinositides, released IP₃ induces conformational change in IP3R2 and increases luciferase activity.

(E) Luminescent kinetics of the NanoBiT-IP₃ sensor after GPCR ligand stimulation. HEK293 cells expressing the sensor alone (Mock) or with a test GPCR (CHRM1) were stimulated with its ligand (acetylcholine, ACh) or vehicle. After ligand addition, a microplate was measured at 10 s interval for 15 min. Each line indicates a luminescent trace of one well, normalized to an initial count, from a representative experiment of at least three independent experiments with similar results.


(F) Validation of $G_{q/11}$ -mediated signal of the NanoBiT-IP₃ sensor. The parental or ΔG_q cells transiently expressing CHRM1 and the NanoBiT-IP₃ sensor were stimulated with vehicle or ACh, and ACh-induced luminescent signal change was normalized to that of vehicle treatment. Bar and error bars represent mean and SEM, respectively, of 5 (parent) or 4 (ΔG_q) independent experiments with each performed in triplicate. *** $p < 0.001$ (t-test).

(G) Measurement of IP₃ formation in EDNRA. The parent, ΔG_q and ΔG_{12} cells transiently expressing EDNRA and the NanoBiT-IP₃ sensor were stimulated with a titrated ligand and ligand-induced luminescent signal was assessed. Symbols and error bars represent mean and SEM of the indicated numbers of independent experiments with each performed in duplicate. Parameters (mean \pm SEM) obtained from the concentration-response curves are shown at the right.

A




B




LogRIA cutoffs ■ -1.0 ■ -0.5 ■ -0.1


C



D



E



F




Figure S7. Predictor Performances, Shortest Path from Contact Network Analysis, DREADD Predictions Scatterplot, Related to Figures 4, 6, and 7

- (A) Radial plot representing Matthew correlation coefficient (MCC) of 5-fold cross validation (averaged over 10 runs).
- (B) Radial plot representing Recall (Sensitivity) of the best performing predictors over the Test set.
- (C) Bar plot representing the recall (sensitivity) of the best performing predictors, trained at different LogRA_i cutoffs, over the test set.
- (D) Example of a shortest communication pathway, depicted on 3D cartoons of the *ADRB2-GNAS* complex (PDB: 3SN6), linking the ligand and G-protein consensus binding pocket pockets.
- (E) Connectivity matrix displaying shortest paths (as intersecting circles) linking residues forming the ligand and G-protein consensus binding pockets (i.e., shown to form such interfaces in at least 50% of the considered structures). Circle color indicates the path length and the diameter is proportional to the number of significant coupling features found at linking positions. Circle rims are red marked if the path is exclusively found on active-like GPCR structures.
- (F) Scatterplot of the relative coupling probabilities of chimeric sequences obtained by swapping on the M3D backbone sequence the sequence stretches corresponding to the ICL3 alone (y axis) or in combination with the C-term (x axis) from the 148 receptors of the chimeric G-protein-based TGF- α shedding assay. The zoomed caption highlights the cluster of chimeric sequences (including GPR183 and GPR132) displaying an increase of coupling probability for GNA12 compared to the reference (i.e., M3D).